



## RESEARCH ARTICLE

F.U. Vet. J. Health Sci.  
2021; 35 (3): 145 - 150  
<http://www.fusabil.org>

### Evaluation of Acute Phase Response and Oxidative Damage in Sheep Naturally Infected with *Streptococcus Pluranimalium* using Haptoglobin, Nitric Oxide, and Malondialdehyde Levels\*

Akın KIRBAŞ<sup>1, a</sup>  
Ramazan KAMAN<sup>2, b</sup>  
Oğuz MERHAN<sup>3, c</sup>  
Kadir BOZUKLUHAN<sup>4, d</sup>  
Şükrü DEĞİRMENÇAY<sup>2, e</sup>  
Muhammed Sertaç EROĞLU<sup>2, f</sup>

<sup>1</sup> Yozgat Bozok University,  
Faculty of Veterinary  
Medicine,  
Department of Internal  
Medicine,  
Yozgat, TURKEY

<sup>2</sup> Atatürk University,  
Faculty of Veterinary  
Medicine,  
Department of Internal  
Medicine,  
Erzurum, TURKEY

<sup>3</sup> Kafkas University,  
Faculty of Veterinary  
Medicine,  
Department of  
Biochemistry,  
Kars, TURKEY

<sup>4</sup> Kafkas University,  
Kars School of Higher  
Vocational Education,  
Department of Veterinary  
Health,  
Kars, TURKEY

<sup>a</sup> ORCID: 0000-0001-9159-3240

<sup>b</sup> ORCID: 0000-0001-6304-6063

<sup>c</sup> ORCID: 0000-0002-3399-0667

<sup>d</sup> ORCID: 0000-0003-4929-5156

<sup>e</sup> ORCID: 0000-0002-3920-6343

<sup>f</sup> ORCID: 0000-0003-1061-8421

Received : 08.09.2021

Accepted : 22.09.2021

#### Correspondence Yazışma Adresi

Akın KIRBAŞ

Yozgat Bozok University,  
Faculty of Veterinary  
Medicine,  
Department of Internal  
Medicine,  
Yozgat – TURKEY

[akindahiliye55@yahoo.com](mailto:akindahiliye55@yahoo.com)

The purpose of this study was to evaluate of acute phase response and oxidative damage in sheep naturally infected with *Streptococcus pluranimalium* using haptoglobin (Hp), nitric oxide (NO), and malondialdehyde (MDA) levels. The animal material of the study consisted of 60 sheep infected with *S. pluranimalium* (Infected group) aged between 4-6 years and 15 healthy sheep (Control group) aged 4-6 years old. Skin scrapings samples were obtained from infected sheep for microbiological analyses. Blood samples were taken from *vena jugularis* of animals in to vacuum tubes for biochemical analysis. From the samples taken from infected sheep, *S. pluranimalium* was identified by using VITEK 2 Compact Bacterial Identification and Monitoring System. Serum Hp and NO levels were determined by the spectrophotometric method and MDA levels were determined by the thiobarbituric acid (TBA) reaction. Hp ( $0.17 \pm 0.01$  g/L), ( $P < 0.001$ ) MDA ( $1.67 \pm 0.04$   $\mu$ mol/L) and NO ( $31.14 \pm 1.61$   $\mu$ mol/L) ( $P < 0.01$ ) levels of the infected group were found to be significantly higher than the control group. It was concluded that high serum MDA and NO levels indicate that oxidative damage takes place in naturally infected sheep with *S. pluranimalium*. However, it can be assumed that higher Hp and low Alb concentrations might be an indicator of acute-phase response in the infected animals.

**Key Words:** Acute phase response, oxidative damage, sheep, *Streptococcus pluranimalium*

#### *Streptococcus Pluranimalium* ile Doğal Enfekte Koyunlarda Akut Faz Yanıt ve Oksidatif Hasarın Haptoglobin, Nitrik Oksit ve Malondialdehit Düzeyleri Kullanılarak Değerlendirilmesi

Bu çalışmanın amacı *Streptococcus pluranimalium* ile doğal enfekte koyunlarda haptoglobin (Hp), nitrik oksit (NO) ve malondialdehit (MDA) düzeylerini kullanarak akut faz yanıtı ve oksidatif hasarı değerlendirmektir. Çalışmanın hayvan materyalini 4-6 yaş arasında olan *S. pluranimalium* ile enfekte 60 adet koyun (Enfekte grup) ve 4-6 yaş arasında olan 15 adet sağlıklı koyun (Kontrol grubu) oluşturmuştur. Mikrobiyolojik analizler için enfekte koyunlardan deri kazıntısı ve biyokimyasal analizler için hayvanların vena jugularislerinden kan örnekleri alındı. Enfekte koyunlardan alınan örneklerden *S. pluranimalium*, VITEK 2 Kompakt Bakteri Tanımlama ve İzleme Sistemi kullanılarak tanımlandı. Serum Hp ve NO düzeyleri spektrofotometrik yöntemle, MDA düzeyleri ise tiyobarbitürik asit (TBA) reaksiyonu ile belirlendi. Enfekte grubun Hp ( $0.17 \pm 0.01$  g/L), ( $P < 0.001$ ) MDA ( $1.67 \pm 0.04$   $\mu$ mol/L) ve NO ( $31.14 \pm 1.61$   $\mu$ mol/L) ( $P < 0.01$ ) düzeyleri kontrol grubundan anlamlı olarak daha yüksek bulundu. Sonuç olarak, yüksek serum MDA ve NO düzeylerinin *S. pluranimalium* ile doğal enfekte koyunlarda oksidatif hasarın meydana geldiğini gösterdiği sonucuna varıldı. Bununla birlikte, daha yüksek Hp ve düşük Alb konsantrasyonlarının, enfekte hayvanlarda akut faz yanıtının bir göstergesi olabileceği varsayılabilir.

**Anahtar Kelimeler:** Akut faz yanıt, oksidatif hasar, koyun, *Streptococcus pluranimalium*

#### Introduction

*Streptococcus pluranimalium* was first described as a new species of the *Streptococcus* genus in 1999 (1). *S. pluranimalium* is promiscuous, in terms of its host and tissue tropism since it has been isolated from various tissues of multiple domestic animals and humans (1). *S. pluranimalium* has been reported to cause subclinical mastitis in dairy cows (1, 2), and many bovine reproductive diseases (abortion, stillbirth, metritis vaginitis, and vulvitis) (2, 3). Additionally, this bacterium has been isolated from human patients with subdural empyema, infective endocarditis, and brain abscess (4-6). On the other hand; pathogenic mechanisms of *S. pluranimalium* are actually unknown at the present time. However, there are insufficient reported data on *S. pluranimalium* infection in sheep.

Reactive oxygen species (ROS) are produced during physiological and metabolic processes and can lead to detrimental oxidative reactions under conditions of excessive production (7, 8). Oxidative damage may develop during situations such as protozoal (9, 10), viral (11, 12) and bacterial infections (13). The primary targets of ROS are

\* 2<sup>nd</sup> International and 13<sup>th</sup> National Veterinary Internal Medicine Congress, 11–13 October 2019, Ankara/Turkey.

polyunsaturated fatty acids (PUFAs) in cell membranes. The resulting lipid peroxidation (LPO) may lead to the damage of the cell structure and functions (14, 15). In addition, the decomposition of lipid hydroperoxides yields a wide variety of end-products, including malondialdehyde (MDA). One of the most frequently used ROS biomarkers, providing an indication of the overall LPO intensity is MDA (13-16).

Nitric oxide (NO) is a signalling molecule that plays a key role in the pathogenesis of inflammation (17). Its release is stimulated by macrophages activated by cytokines (18). In conclusion, large amounts of NO are synthesized, exceeding the physiological NO production (17, 18). NO is a substantial molecule included in physiological and pathological processes in animals (17). The activity of NO in cellular defense mechanisms includes participation in tissue injury and the mediation of inflammatory processes and apoptosis (18). However, NO is stated to play an important role in the primary defense against several species of bacteria (19, 20), viruses (11, 21, 22) and parasites (23).

Acute-phase response (APR) mostly occurs during infection and inflammation. The aim of these reactions is to isolate and destroy the infectious agents to prevent ongoing tissue damage, and to restore homeostasis. One of the main features of APR is the hepatic production of acute-phase proteins (APPs). The blood concentration of APPs generally increases within 8h of stimulation, reaches the maximum level in 24-48h (24). APPs may be used for the differentiation of bacterial and viral infections, for the differential diagnosis of clinical, subclinical, acute and chronic diseases (24, 25). The most important major APPs for the determination of sheep diseases are haptoglobin (Hp) (24-26). Hp is APP binding free haemoglobin in the blood. Hp concentrations are increased during acute infection but decreased with treatment or chronicity (24, 25). Nevertheless, their concentration remains high in chronic cases if stimulation continues (25). However, no study was encountered in sheep on the related infection, and no adequate studies have been found on oxidative damage and acute phase response. Therefore, in this study, it was aimed to evaluate the acute phase response and oxidative damage in sheep naturally infected with *S.pluranimalium* using Hp, NO, and MDA levels.

## Material and Methods

**Research and Publication Ethics:** On a farm located in Erzurum with 150 Morkaraman sheep aged 4-6 years old, the owner applied us for diagnosis and treatment as some sheep had pyrexia, anorexia, excessive weakening, necrotic ulcerative dermatitis in the tail area, loss of appetite and weight loss. The flock of sheep was examined after obtaining the "informed consent form" from the owner. The animal material of this study consisted of 60 symptomatic sheep (infected group) and 15 healthy sheep (control group).

**Animals:** The animal material of the study consisted of infected sheep from a 150-animal sheep

herd (Morkaraman breed) in Erzurum province and healthy sheep from the same herd. The infected group consisted of 60 infected sheep aged 4-6 years (Morkaraman breed); while the control group consisted of 15 healthy sheep (Morkaraman breed) aged 4-6 years old. This study was conducted in accordance with ethical rules.

**Blood and Skin Scrapings Sampling:** Blood samples were taken from the *jugular veins* into vacuum tubes without anticoagulant (Vacutainer, BD-Plymouth, UK) for serum analyses. The blood samples were centrifuged at 3000 g for 10 min at room temperature. The sera were separated and stored at -80°C until analysed. Skin scrapings samples were obtained from sick sheep with suspected infection for microbiological analysis.

**Bacteriological Analysis:** From the samples taken from infected sheep, *S. pluranimalium* was isolated and identified by using VITEK 2 Compact Bacterial Identification and Monitoring System (Biomérieux, Inc., Hazelwood, MO, USA).

## Biochemical Analyses

**Malondialdehyde (MDA) and Nitric Oxide (NO) analyses:** Serum MDA levels were measured by the thiobarbituric acid (TBA) reaction according to the method described by Yoshioka et al. (27). The end products were read at 535 nm. Serum NO levels were determined according to the method described by Miranda et al. (28). Nitrate is reduced to nitrite by  $\text{VCl}_3$ , and then in an acidic environment nitrite was reacted with sulphanilamide to produce coloured diazonium compound, which was read at 540 nm.

**Haptoglobin (Hp) and Albumin (Alb) Analyses:** Serum Hp concentrations were determined spectrophotometrically according to the method which have been previously reported by Skinner et al. (29). Serum Alb concentrations also were measured spectrophotometrically (Epoch, Biotek, USA) using a commercial test kit (Biolabo, France).

**Statistical Analysis:** The SPSS software program (Version 20.0, SPSS Inc., Chicago, IL, USA) was used for statistical analysis. The distribution of the data between the groups was evaluated using the Kolmogorov-Smirnov test. Levene's test was used to test whether variances were homogenous. Parametrically distributed groups were compared using the t-test (Independent-Samples t-test). The correlation among parameters was determined by the Pearson Correlation test. All data were presented as the mean and standard error of the mean (Mean $\pm$ SEM). The results were assessed at a 95% confidence interval and a significance level of  $P < 0.05$  (30).

## Results

Pyrexia, anorexia, excessive weakening, necrotic ulcerative dermatitis in the tail area, death within 5 to 10 days were important clinical findings in infected sheep (Figure 1).

Serum levels of Hp, Alb, MDA and NO in healthy and infected sheep are given in Table 1. Hp ( $0.17 \pm 0.01$  g/L) ( $P < 0.001$ ), MDA ( $1.67 \pm 0.04$   $\mu\text{mol/L}$ ) and NO ( $31.14 \pm 1.61$   $\mu\text{mol/L}$ ) ( $P < 0.01$ ) levels of the infected group were found significantly higher than the control group. The serum Alb concentrations in the infected group were not statistically different from the control group ( $P > 0.05$ ). Correlations between parameters of the infected group were presented in Table 2. While there was a positive correlation between Hp, NO, and MDA levels of the infected group, there was a significant positive correlation between Hp concentrations and MDA levels ( $P < 0.05$ ). However, a negative correlation was determined between NO, MDA levels, and Alb concentrations.



**Figure 1.** Necrotic-ulcerative dermatitis in the ventral of the tail, ranging from 1-3 cm to 10-15 cm in size. (A): Primary effect (bold arrow), (B-C): Progressing lesions, ulcerations (thin arrow), necrotic mass (arrow heads).

**Table 1.** Hp, Alb, MDA and NO levels in healthy and infected sheep

| Parameters                | Healthy<br>(n=15) | Infected<br>(n=60) | P         |
|---------------------------|-------------------|--------------------|-----------|
|                           | Mean $\pm$ SEM    | Mean $\pm$ SEM     |           |
| Hp (g/L)                  | $0.11 \pm 0.01$   | $0.17 \pm 0.01$    | $< 0.001$ |
| Alb (g/dL)                | $3.18 \pm 0.12$   | $2.92 \pm 0.06$    | $> 0.05$  |
| MDA ( $\mu\text{mol/L}$ ) | $1.35 \pm 0.10$   | $1.67 \pm 0.04$    | $< 0.01$  |
| NO ( $\mu\text{mol/L}$ )  | $22.07 \pm 3.70$  | $31.14 \pm 1.61$   | $< 0.01$  |

Hp: haptoglobin; Alb: albumin; MDA: malondialdehyde;  
NO: nitric oxide

**Table 2.** Correlation between parameters in infected sheep

| Parameters | Hp    | NO    | MDA    | Alb    |
|------------|-------|-------|--------|--------|
| Hp         | 1.000 | 0.198 | 0.243* | 0.012  |
| NO         |       | 1.000 | 0.129  | -0.141 |
| MDA        |       |       | 1.000  | -0.174 |
| Alb        |       |       |        | 1.000  |

Hp: haptoglobin; NO: nitric oxide; MDA: malondialdehyde;  
Alb: albumin; \*:  $P < 0.05$

## Discussion

There have been insufficient studies of *S. pluranimalium* infection in sheep worldwide. Therefore, this study aimed to evaluate the acute-phase response and oxidative status in infected sheep. While this bacterium causes problems with endocarditis and brain abscess in humans (4-6), it is reported that it causes mastitis, abortion, and stillbirths in cattle (2, 3),

meningoventriculitis in a calf (31), respiratory disease in dogs (32), and septicaemia and endocarditis in broiler chickens (33). On the other hand, in only one report, Foster et al. (34) isolated the agent from two sheep abortion materials, especially from stomach and liver samples of the fetus. At the same time, as clinical findings, it was reported that the sheep were pyrexia at the time of the abortion. In the presented study, pyrexia, anorexia, excessive weakening, weight loss, necrotic ulcerative dermatitis in the tail area (Figure 1), and death within 5 to 10 days were important clinical findings in the infected sheep.

Common causes of oxidative stress are infections, inflammation, and toxemia (35). Infections and inflammations activate inflammatory cells that play important roles in the host's defense (15). Since major inflammatory cells, such as neutrophils and/or macrophages, generate a variety of ROS and release various proteases, tissue damage and destruction occur (15, 35). In addition, one of the many biological targets of oxidative stress is lipids. PUFAs, in particular, are the frequently targeted class of biomolecules. ROS-induced oxidation of PUFAs in biological systems results in the formation of LPO products (15, 35). MDA is a breakdown product that is quantified as a measure of lipid hydroperoxides, and is accepted as an indicator of elevated oxidative stress in the body (36, 37). In sheep, oxidant status has been identified in metabolic (38, 39), parasitic (9, 10, 40), and viral diseases (11, 12), as well as in physiological conditions, such as age (41) and pregnancy (42). In the present study, serum MDA levels were significantly increased in infected sheep compared to healthy sheep ( $P < 0.01$ ). In this case, it has been consistent with the previous reports. These increases can be considered as an indicator of excessive free radical production in infected sheep.

NO is a prominent molecule in defense against many microorganisms (17, 18). NO, which is produced by macrophages in bacterial infections, shows antibacterial properties against bacteria and host defense depends on the concentration of NO (18). It was reported that NO concentrations in animals with bacterial (19, 20), viral (11, 21, 22) and parasitic diseases (23) increased compared to healthy controls. In addition, it has been determined that NO levels increase in bacterial diseases such as traumatic reticuloperitonitis (43) and traumatic pericarditis (44). Similarly, in the present study, NO concentrations were high in infected sheep as in the above reports. This increase was thought due to inducing of NO synthesis by macrophages defending the organism against infection.

Haptoglobin belongs to a group of transporter plasma proteins that bind the free haemoglobin (45). This defines its bacteriostatic effect as well as antioxidant activity (26, 45). Many studies have demonstrated the importance of Hp as a clinically helpful parameter for measuring the occurrence and severity of inflammatory responses in sheep with various infectious diseases (12, 46-52).

Pepin et al. (46) investigated the changes in Hp concentrations in lambs experimentally infected by *Corynebacterium pseudotuberculosis*. They reported the rapidly increasing Hp concentration to peak in plasma after subcutaneous inoculation. In a similar study, Eckersall et al. (47) explored the differences in plasma levels of Hp during experimental ovine caseous lymphadenitis induced by *C. pseudotuberculosis*. The results showed significantly raised Hp concentrations on day 7 after inoculation and these values were statistically significant until the 15th day. In another study, Bastos et al. (49) also stated that there was no significant difference between serum Hp concentrations in seropositive and seronegative sheep during caseous lymphadenitis in Santa-Ines sheep. However, later in 11 sheep that have not developed peripheral abscesses, a significantly higher Hp concentration was observed. Fasulkov et al. (50) investigated plasma Hp concentrations during experimentally induced *Staphylococcus aureus* mastitis in goats and showed an increase of the Hp as early as the 8th hour with the most significant differences from baseline values by the 24th and 48th h after infection. On the other hand, Chalmeh et al. (51) found a rapid elevation in Hp levels during experimentally induced endotoxaemia in sheep by lipopolysaccharide from *Escherichia coli*. In the present study, Hp concentrations were significantly higher in infected sheep (Hp;  $0.17 \pm 0.01$ g/L versus  $0.11 \pm 0.01$ g/L) compared to the healthy control group ( $P < 0.001$ ). These values were observed to be similar to different studies

conducted in previous years (45-49). Therefore, it can be said that Hp concentrations are important in the pathogenesis and monitoring of *S. pluranimalium* infection. In addition, there was a significantly positive correlation between Hp and MDA levels of the infected sheep ( $P < 0.05$ ). These findings may indicate that APR and oxidative damage occurs simultaneously in infected sheep and triggers each other.

Albumin is the major negative APP. During the APR the demand for amino acids for synthesis of the positive APPs is markedly increased, which necessitates reprioritization of hepatic protein synthesis. Therefore, Alb synthesis is down-regulated and amino acids are shunted into the synthesis of positive APPs (53, 54). In addition, it has been stated that the serum concentrations of Alb, are affected by impaired liver function, reduced intestinal absorption, and starvation (24, 55). In the presented study, the serum Alb concentrations in infected sheep were not statistically different from the control group. However, the decrease in Alb concentration may be related to the synthesis of APPs and this finding may indicate that hepatic Alb synthesis was affected by APR.

In conclusion, high serum MDA and NO levels indicate that oxidative stress takes place in naturally infected sheep with *S. pluranimalium*. Higher Hp and low Alb concentrations in the infected animals compared to the healthy ones is also an indicator of APR in these animals.

## References

- Devriese LA, Vandamme P, Collins MD, et al. *Streptococcus pluranimalium* sp. nov., from cattle and other animals. Int J Syst Bacteriol 1999; 49: 1221-1226.
- Twomey DF, Carson T, Foster G, Koylass MS, Whatmore AM. Phenotypic characterization and 16S rRNA sequence analysis of veterinary isolates of *Streptococcus pluranimalium*. Vet J 2012; 192: 236-238.
- Foster G, Barley J, Howie F, et al. *Streptococcus pluranimalium* in bovine reproductive disease. Vet Rec 2008; 163: 638.
- Aryasinghe L, Sabbar S, Kazim Y, Awan LM, Khan HK. *Streptococcus pluranimalium*: A novel human pathogen? Int J Surg Case Rep 2014; 5: 1242-1246.
- Fotoglidis A, Pagourelas E, Kyriakou P, Vassilikos V. Endocarditis caused by unusual *Streptococcus species* (*Streptococcus pluranimalium*). Hippokratia 2015; 19: 182-185.
- Maher G, Beniwal M, Bahubali V, et al. *Streptococcus pluranimalium*: An emerging animal streptococcal species as a causative agent of human brain abscess. World Neurosurg 2018; 115: 208-212.
- Celi P. Biomarkers of oxidative stress in ruminant medicine. Immunopharmacol and Toxicol 2011; 33: 233-240.
- Lykkesfeldt J, Svendsen O. Oxidants and antioxidants in disease: Oxidative stress in farm animals. Vet J 2007; 173: 502-511.
- Nazifi S, Razavi SM, Kianiamin P, Rakhshandehroo E. Evaluation of erythrocyte antioxidants mechanisms: Antioxidants enzymes, lipid peroxidation, and serum trace elements associated with progressive anemia in ovine malignant theileriosis. Parasitol Res 2011; 109: 275-281.
- Esmailnejad B, Tavassoli M, Asri-Rezaei S, Dalir-Naghadeh B. Evaluation of antioxidant status and oxidative stress in sheep naturally infected with *Babesia ovis*. Vet Parasitol 2012; 185: 124-130.
- Kirmizigul AH, Ogun M, Ozen H, et al. Oxidative stress and total sialic acid levels in sheep naturally infected with pox virus. Pak Vet J 2016; 36: 312-315.
- Bozukluhan K, Merhan O, Gokce HI, et al. Determination of some acute phase proteins, biochemical parameters and oxidative stress in sheep with naturally infected sheepox virus. Kafkas Univ Vet Fak Derg 2018; 24: 437-441.
- Cenesiz S. The Role of Oxidant and Antioxidant Parameters in the Infectious Diseases: A Systematic Literature Review. Kafkas Univ Vet Fak Derg 2020; 26: 849-858.
- Halliwell B, Chirico S. Lipid peroxidation: Its mechanism, measurement, and significance. Am J Clin Nutr 1993; 57: 715-725.
- Winrow VR, Winyard PG, Morris CJ, Blake DR. Free radicals in inflammation: Second messengers and mediators of tissue destruction. Br Med Bull 1993; 49: 506-522.

16. Palmieri B, Sblendorio V. Oxidative stress tests: Overview on reability and use part I. Eur Rev Med Pharmacol Sci 2007; 11: 309-342.
17. Sharma JN, Al-Omran A, Parvathy SS. Role of nitric oxide in inflammatory diseases. Inflammopharmacol 2007; 15: 252-259.
18. Wallace JL. Nitric oxide as a regulator of inflammatory processes. Mem Inst Oswaldo Cruz 2005; 100 (Suppl. I): 5-9.
19. Nisbet C, Yarim GF, Ciftci A, Cenesiz S, Ciftci G. Investigation of serum nitric oxide and malondialdehyde levels in cattle infected with *Brucella abortus*. Ankara Univ Vet Fak Derg 2007; 54: 159-163.
20. Hanedan B, Kirbas A, Kandemir FM, Aktas MS, Yildiz A. Evaluation of arginase activity, nitric oxide, and oxidative stress status in sheep with Contagious agalactiae. Acta Vet Hung 2017; 65: 394-401.
21. Kandemir FM, Issi M, Benzer F, et al. Plasma nitric oxide concentrations and erythrocyte arginase activities in lambs with contagious ecthyma. Revue Méd Vét 2011; 162: 275-278.
22. Aslan O, Gencay Goksu A, Apaydin N. The evaluation of oxidative stress in lambs with Pestivirus infection. J Hellenic Vet Med Soc 2017; 68: 299-306.
23. Kontas T, Salmanoglu B. Tumour necrosis factor- $\alpha$ , adenosine deaminase and nitric oxide levels in cattle babesiosis before and after treatment. Bull Vet Inst Pulawy 2006; 50: 485-487.
24. Gruys E, Toussiant M, Niewald TA. Acute phase reaction and acute phase proteins. J Zhejiang Univ Sci 2005; 11: 1045-1056.
25. Petersen HH, Nielsen JP, Heegard PMH. Application of acute phase protein measurements in veterinary clinical chemistry. Vet Res 2004; 35: 163-167.
26. Cecilian F, Ceron JJ, Eckersall PD, Sauerwein H. Acute phase proteins in ruminants. J Proteom 2012; 275: 4207-4231.
27. Yoshioka T, Kawada K, Shimada T, Mori M. Lipid peroxidation in maternal and cord blood and protective mechanism against activated-oxygen toxicity in the blood. Am J Obstet Gynecol 1979; 135: 372-376.
28. Miranda KM, Espey MG, Wink DA. A rapid, simple spectrophotometric method for simultaneous detection of nitrate and nitrite. Nitric Oxide 2001; 5: 62-71.
29. Skinner JG, Brown RA, Roberts L. Bovine haptoglobin response in clinically defined field conditions. Vet Rec 1991; 128:147-149.
30. Ozdamar K. SPSS Applied Basic Statistics for Education Health and Social Sciences. Eskisehir: Nisan Bookstore, 2018.
31. Seimiya YM, Takahashi M, Kudo T, Sasaki K. Meningoventriculitis caused by *Streptococcus pluranimalium* in a neonatal calf of premature birth. J Vet Med Sci 2007; 69: 657-660.
32. Kalhor DH, Luo S, Xie X, et al. *Streptococcus pluranimalium* isolated from a canine respiratory case: Identification and experimental infection in mice. Pak Vet J 2015; 35: 388-390.
33. Hedegaard L, Christensen H, Chadfield MS, Christensen JP, Bisgaard M. Association of *Streptococcus pluranimalium* with valvular endocarditis and septicemia in adult broiler parents. Avian Pathol 2009; 38:155-160.
34. Foster G, Hunter L, Baird G, Koylass MS, Whatmore AM. *Streptococcus pluranimalium* in ovine reproductive material. Vet Rec 2010; 166: 246.
35. Mandelker L. Introduction to oxidative stress and mitochondrial dysfunction. Vet Clin Small Anim Pract 2008; 38: 1-30.
36. Singh Z, Karthigesu IP, Singh P, Kaur R. Use of malondialdehyde as a biomarker for assessing oxidative stress in different disease pathologies: A review. Iranian J Publ Health 2014; 43 (Suppl.3): 7-16.
37. Tsikas D. Assessment of lipid peroxidation by measuring malondialdehyde (MDA) and relatives in biological samples: Analytical and biological challenges. Anal Biochem 2017; 524: 13-30.
38. Al-Qudah KM. Oxidant and antioxidant profile of hyperketonemic ewes affected by pregnancy toxemia. Vet Clin Pathol 2011; 40: 60-65.
39. Kirbas A, Yildirim BA, Baydar E, Kandemir FM. Status of lipid peroxidation and some antioxidants in sheep with acute ruminal lactic acidosis. Med Weter 2014; 70: 357-361.
40. Aktas MS, Kandemir FM, Kirbas A, Hanedan B, Aydin MA. Evaluation of oxidative stress in sheep infected with *Psoroptes ovis* using total antioxidant capacity, total oxidant status, and malondialdehyde level. J Vet Res 2017; 61: 197-201.
41. Amoli JS, Baghbanzadeh A. Oxidative stress in Sahal sheep of different age groups. Turk J Vet Anim Sci 2010; 34: 379-383.
42. Erisir M, Benzer F, Kandemir FM. Changes in the rate of lipid peroxidation in plasma and selected blood antioxidants before and during pregnancy in ewes. Acta Vet Brno 2009; 78: 237-242.
43. Kirbas A, Baydar E, Kandemir FM. Assessment of plasma nitric oxide concentration and erythrocyte arginase activity in dairy cows with traumatic reticuloperitonitis. J Hellenic Vet Med Soc 2019; 70: 1833-1840.
44. Ozkan C, Altug N, Kaya A, Basbugan Y. Serum nitric oxide concentrations in cattle with traumatic pericarditis. Van Vet J 2012; 23: 131-135.
45. Tothova C, Nagy O, Kovac G. Acute phase proteins and their use in the diagnosis of diseases in ruminants: A review. Vet Med 2014; 59: 163-180.
46. Pepin M, Pardon P, Lantier F, et al. Experimental *Corynebacterium pseudotuberculosis* infection in lambs: Kinetics of bacterial dissemination and inflammation. Vet Microbiol 1991; 26: 381-392.
47. Eckersall PD, Lawson F, Bence L, et al. Acute phase protein response in an experimental model of ovine caseous lymphadenitis. BMC Vet Res 2007; 3: 1-6.
48. Eckersall PD, Bell R. Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine. Vet J 2010; 185: 23-27.

49. Bastos BL, Meyer R, Guimarães JE, et al. Haptoglobin and fibrinogen concentrations and leukocyte counts in the clinical investigation of caseous lymphadenitis in sheep. *Vet Clin Pathol* 2011; 40: 496-503.
50. Fasulkov I, Karadaev M, Vasilev N, Urumova V, Mircheva T. Determination of plasma fibrinogen and haptoglobin, hematological and blood biochemical changes in Bulgarian local goats with experimentally induced *Staphylococcus aureus* mastitis. *Turk J Vet Anim Sci* 2014; 38: 439-444.
51. Chalmeh A, Badiei K, Pourjafar M, Nazifi S. Acute phase response in experimentally *Escherichia coli* serotype O55:B5 induced endotoxemia and its comparative treatment with dexamethasone and flunixin meglumine in Iranian fat-tailed sheep. *Vet Arhiv* 2013; 83: 301-312.
52. Iliev PT, Georgieva TM. Acute phase biomarkers of diseases in small ruminants: an overview. *Bulg J Vet Med* 2019; 22: 1-12.
53. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *N Engl J Med* 1999; 340: 448-454.
54. Kirbas A, Ozkanlar Y, Aktas MS, et al. Acute phase biomarkers for inflammatory response in dairy cows with traumatic reticuloperitonitis. *Isr J Vet Med* 2015; 70: 23-29.
55. Kaneko JJ, Harvey JW, Bruss ML. *Clinical Biochemistry of Domestic Animals*. 6th Edition, New York: Academic Press, 2008.