

## Subepiglottic Cyst in A Neonatal Calf, in Turkey

Ali Haydar KIRMIZIGÜL\*   
Erhan GÖKÇE\*

Özgür AKSOY\*\*  
Başak KURT\*\*

İsa ÖZAYDIN\*\*  
Celal Şahin ERMUTLU\*\*

\* Department of Internal Medicine, Faculty of Veterinary Medicine, Kafkas University, Kars - TURKEY

\*\* Department of Surgery, Faculty of Veterinary Medicine, Kafkas University, Kars - TURKEY

Yayın Kodu (Article Code): 2008/51-E

Dear Editor,

We would like to report a rare case of subepiglottic cyst in animals and is reported for the first time in a neonatal calf in our country.

Subepiglottic cysts are rare condition in human beings <sup>1-3</sup>, horses <sup>4,5</sup>, dogs <sup>6</sup> and cats <sup>7</sup>. Only one case of subepiglottic cysts has been reported in cattle <sup>8</sup>. Inspiratory stridor and varying degrees of upper airway obstruction are the most frequently clinical findings in subepiglottic cyst. Other common findings included feeding difficulties and episodes of cyanosis <sup>9-11</sup>.

A 25 day old Simmental calf with respiratory distress was admitted to clinic of Department of Internal Medicine, Faculty of Veterinary Medicine, University of Kafkas. Case history revealed that the calf had respiratory distress since birth and had been treated by a veterinarian but did not improved in spite of fairly good general condition. Clinical examination of the calf disclosed severe respiratory distress, cyanosis of mucous membranes, hyperaemia in conjunctivas, increased respiratory (50 breaths/min) and heart rates (140 beats/min). Palpation of laryngeal area resulted in severe respiratory distress and cessation of respiration for short while. Endotracheal tube was placed in to relieve respiratory distress. Following local anaesthesia for tracheotomy animal died in a short while. Necropsy of subepiglottic area revealed a cyst of 3 cm in diameter in left ventrolateral larynx (*Figure 1*) filled with grey-whitish fluid.

As a result, this was the first case of subepiglottic cyst determined in a calf in our country and subepiglottic cyst should also be considered in cases with upper respiratory tracts.

### REFERENCES

1. **Berger G, Averbuch E, Zilka K, Berger R, Ophir D:** Adult vallecular cyst: Thirteen-year experience. *Otolaryngology -Head and Neck Surgery*, 138, 321-327, 2008.
2. **Righini CA, Kadaoui H, Morel N, Llerena C, Reyt E:** Stridor in a newborn caused by a congenital laryngeal saccular cyst. *Int j Pediatr Otorhinolaryngol*, 1, 145-149, 2006.
3. **Rivo J, Matot I:** Asymptomatic vallecular cyst: Airway management considerations. *J Clin Anesth*, 13, 383-386, 2001.
4. **Hay WP, Baskett A, Abdy MJ:** Complete upper airway obstruction and syncope caused by a subepiglottic cyst in a horse. *Equine Vet J*, 29, 75-76, 1997.
5. **Stick JA, Boles C:** Subepiglottic cyst in three foals. *J Am Vet Med Assoc*, 177, 62-64, 1980.
6. **Caywood D, Wallace LJ, Alsaker RD, Lord PF:** Laryngeal cyst in dog - case-report. *J Am Anim Hosp Assoc*, 13, 87-91, 1977.
7. **Rudolf H, Lane JG, Brown PJ, Mackay A:** Ultrasonographic diagnosis of a laryngeal cyst in a cat. *J Small Anim Pract*, 40, 275-277, 1999.
8. **Mattoon JS, Andrews D, Jones SL, Linford RL:** Subepiglottic cyst causing upper airway obstruction in a neonatal calf. *J Am Vet Med Assoc*, 199, 747-749, 1991.
9. **Gutierrez JP, Berkowitz RG, Robertson CF:** Vallecular cysts in newborns and young infants, *Pediatr Pulmonol*, 27, 282-285, 1999.
10. **Hsieh WS, Yang PH, Wong KS, Li HY, Wang ECR, Yeh TF:** *Eur J Pediatr*, 159, 79-81, 2000.
11. **Gluckman PGC, Chu TWF, van Hasselt CA:** Neonatal vallecular cysts and failure to thrive, *J Laryngol Otol*, 106, 448-449, 1992.




İletişim (Correspondence)



+90 474 2426800/1251



ahkirmizigul@hotmail.com



**Fig 1.** View of subepiglottic area with a cyst of 3 cm in diameter in left ventrolateral larynges

**Şekil 1.** Subepiglottik bölgede, larinksin sol ventrolateralinde yaklaşık 3 cm çapında kistin görüntüsü