

The Change in Luteal Blood Flow and Luteal Size after Beta Carotene and GnRH Injections in Early Pregnant Dairy Cows

Serhan Serhat AY ¹ İbrahim KÜÇÜKASLAN ² Duygu KAYA ³ Serkan Barış MÜLAZIMOĞLU ⁴ Birten EMRE ⁵ Cihan KAÇAR ³ Hakan KALENDER ⁶ Murat FINDIK ¹ Henrich BOLLWEIN ⁷ Martin RIEGLER ⁸ Sabine SCHÄFER-SOMI ⁸ Jakop SCHOLBACH ⁹ Selim ASLAN ⁴

- ¹ Ondokuz Mayıs University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-55139 Samsun - TURKEY
² Dicle University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-21280 Diyarbakır - TURKEY
³ Kafkas University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-36040 Kars - TURKEY
⁴ Ankara University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-06110 Ankara - TURKEY
⁵ Harran University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-63200 Şanlıurfa - TURKEY
⁶ Kırıkkale University, Department of Obstetrics and Gynecology, Faculty of Veterinary Medicine, TR-71451 Kırıkkale - TURKEY
⁷ Clinic for Cattle, School of Veterinary Medicine Hannover, D-30559 Hannover - GERMANY
⁸ Centre for Artificial Insemination and Embryo Transfer, University of Veterinary Science, A-1210 Vienna - AUSTRIA
⁹ Chameleon Software, D-04105 Leipzig - GERMANY

Makale Kodu (Article Code): KVFD-2012-6959

Summary

The aim of this study was to determine the effectiveness of intramuscular injections of beta-carotene (β C) and GnRH on luteal size (LS), luteal blood flow (LBF) and serum β C concentrations in early pregnant cows. Twenty-nine Holstein-Friesian cows with a mature corpus luteum (>19 mm) were randomly assigned to two groups: β C not received (β C-; $n=15$) or received (β C+; $n=14$). All cows were treated with PGF₂ α and inseminated twice, 48 and 72h after the treatment. Last AI was considered to be day 0. All cows received GnRH on day inseminations, 7 and 17. Different from the β C-, the β C+ group received β C intramuscularly on day 7 and 17. In both groups, measurement of LS and LBF were performed on days 7, 10, 17, 27 and 37 by transrectal B-mode and colour Doppler ultrasonography. Blood samples were collected on each examination day. Only cows that became pregnant were included in the statistical evaluation. The concentration of β C in the β C+ group was higher than in the β C- at all examination days except day 17 ($P<0.05$). There was no significant difference between groups concerning the progesterone concentrations ($P>0.05$). The LS and LBF of β C+ group on day 7 ($P<0.05$) and 27 ($P<0.01$) was higher than in the β C- group and values increased significantly until day 37 (LS: $P<0.05$, LBF: $P<0.01$). We conclude that β C injections significantly increased serum β C concentrations, as well as LS and LBF.

Keywords: Beta carotene, Dairy cows, Luteal blood flow, Luteal size

Erken Gebe Sütçü İneklerde Beta Karoten ve GnRH Enjeksiyonlarından Sonra Luteal Kan Akışı ve Luteal Büyüklükteki Değişiklikler

Özet

Sunulan çalışmada erken gebe ineklere kas içi yapılan beta-karoten (β C) ve GnRH enjeksiyonlarının luteal büyüklük (LS), luteal kan akışkanlığı (LBF) ve serum β C düzeyine etkilerinin belirlenmesi amaçlandı. Olgun korpus luteuma (>19 mm) sahip toplam 29 Holstein-Friesian inek rastgele olarak β C uygulanmayan (β C-; $n=15$) ve uygulanan (β C+; $n=14$) olmak üzere iki gruba ayrıldı. İneklerin hepsi PGF₂ α uygulamasından 48 ve 72 saat sonra tohumlandı. Son tohumlama günü 0. gün olarak kabul edildi. Tohumlamalar sırasında, ayrıca 7. ve 17. günde tüm hayvanlara GnRH uygulandı. β C- gruptan farklı olarak β C+ gruba 7 ve 17. günde β C kas içi uygulandı. Grupların LS ve LBF ölçümleri 7, 10, 17, 27 ve 37. günlerde rektal yolla B-mod ve renkli Doppler ultrasonografi ile gerçekleştirildi. Her bir uygulama gününde kan örnekleri toplandı. Tohumlamalar sonucunda gebe olan inekler istatistiksel değerlendirmede kullanıldı. Serum β C düzeyi β C+ grupta, 17. gün hariç, β C- gruptan yüksek bulundu ($P<0.05$). Grupların progesteron düzeyi arasında ise fark bulunmadı ($P>0.05$). LS ve LBF değerlerinin β C+ grupta 7. ($P<0.05$) ve 27. günlerde ($P<0.01$) β C- gruptan daha yüksek olduğu ve 37. güne kadar önemli düzeyde artarak devam ettiği belirlendi (LS: $P<0.05$, LBF: $P<0.01$). Sonuç olarak β C enjeksiyonlarının serum β C konsantrasyonunu, LS ve LBF'yi önemli derecede yükselttiği belirlendi.

Anahtar sözcükler: Beta karoten, Sütçü inek, Luteal kan akışkanlığı, Luteal büyüklük



İletişim (Correspondence)



+90 362 3121919/1226



serhan.ay@gmail.com; serhan.ay@omu.edu.tr

INTRODUCTION

The corpus luteum (CL) is a transient and steroidogenic organ in the ovary in mammals and its main function is to produce progesterone (P_4), a critical hormone during early pregnancy. It stimulates production of a variety of endometrial secretions that is required for successful embryonic development ¹.

Beta-carotene (β C) is a precursor of vitamin A and has a specific effect on reproductive functions independent from vitamin A ². It has been reported that supplementation of β C improved fertility ³ and decreased puerperal disorders ⁴ in cows. Beta-carotene has an antioxidant effect and bovine CL contains up to more than five times β C from other tissues as liver, adipose tissue or plasma ⁵. This may be necessary due to the free radicals produced during the hydroxylation reactions involved in steroid biosynthesis ⁶. A positive correlation has been reported between β C concentrations of the CL, CL diameter and plasma P_4 concentrations ^{2,7}. Aslan et al. ⁸ reported that delayed formation of the CL and low concentrations of P_4 production after estrus are observed in cows with low serum concentrations of β C. It has been reported that β C increased P_4 secretions by stimulating LH secretion from hypophysis ⁹.

Luteinizing hormone surge was also induced by GnRH treatment ¹⁰. Gümen and Sequin ¹¹ reported a rapid rise in peripheral serum LH levels after the GnRH injections. It is well-known, that the use of GnRH after AI improves pregnancy rates ¹². The use of GnRH at the time of AI or on days 11 to 14 post-AI in cows was found to decrease embryonic loss and to increase pregnancy rate ¹³.

Blood flow in the CL is one of the most important factors for the development of the CL and maintenance of its function ¹⁴, it is also closely associated with the production and release of P_4 ¹⁵. The CL is one of the most highly vascularized organs and receives the greatest rate of blood flow per unit of tissue of any organ in the body ¹⁶. The change in blood flow of CL during the different stages of CL life span was shown by many authors ¹⁷⁻²⁰. Colour Doppler ultrasonography is a useful, non-invasive technique for the evaluation of ovarian vascular function, allowing visual observation of blood flow in a delimited area in the wall of preovulatory follicles, in the CL ¹⁵ or uterus ²¹ of cows.

The aim of this study was to determine the effectiveness of intramuscular injections of β C and GnRH on luteal size (LS), luteal blood flow (LBF) and serum β C concentrations in early pregnant cows.

MATERIAL and METHODS

Animals

Twenty-nine Holstein-Friesian cows aged 2 to 6 years from a commercial dairy farm were used. All animals were lactating for 50 or 80 days prior to the start of the study. The cows were kept in a half-open barn and fed with a total mix ration twice daily that included corn and grass silage, hay, triticale, canola and a balanced grain ration and water intake was ad libitum.

Study Design


Prior to the start of the applications, all cows were controlled for general health status. The body condition score of animals ranged from 3.0 to 3.5 (five point scale). Only cows that were reproductively healthy and had a mature CL (>19 mm) were chosen for the study. Animals were randomly assigned to two groups: beta-carotene not received (β C-; n=15) and beta-carotene received (β C+; n=14). All cows were treated with one dose of $\text{PGF}_{2\alpha}$ (Dinoprost; Enzaprost®, 25 mg/per cow, IM, Ceva-Dif, Turkey) and AI was performed 48 and 72 h after treatment. Last AI was considered to be day 0. A synthetic GnRH analogue (Buserelin; Receptal®, 10 µg/per cow, IM, Intervet, Turkey) was applied during the AIs, as well as on day 7 and 17 in all cows of the study. The cows in the β C+ group additionally received β C (Carofertin®, 200 mg/per cow, IM, Alvetra&Werfft AG, Austria) on the day of the $\text{PGF}_{2\alpha}$ injection, as well as on day 7 and 17 after the last AI. The β C dose was determined in accordance with both previously described ²² and the manufacture's recommendation. Transrectal B-mode and colour Doppler ultrasonography were used to determine and compare LS and LBF on days 7, 10, 17, 27 and 37 in the β C+ and β C- group (Fig. 1). The pregnancy control was performed by transrectal B-mode ultrasonography on d 27.

Ultrasonographic Examinations

Ultrasonographic examinations and image collection were performed as described previously ²³⁻²⁵. The B-mode and colour Doppler mode of the portable LOGIQ Book XP machine (General Electric Healthcare, Solingen, Germany) and its equipment with 10 MHz linear probe were used to measure the LS and LBF. The CL was identified by B-mode, and its image was frozen at the maximum cross-sectional area and stored for further off-line measurements. Colour LBF mapping in various transverse sections was conducted using the power Doppler mode as previously. To minimize the variations in recording, the settings of the Power Doppler

Fig 1. Experimental design of the study (Synch: Synchronization; USG: B-mode Ultrasonography; DP: Colour Doppler Ultrasonography; BS: Blood samples)

Şekil 1. Çalışma planı (Synch: Senkronizasyon; USG: B-mod Ultrasonografi; DP: Renkli Doppler Ultrasonografi; BS: Kan örneği)



system were fixed and used for all examinations. The entire cross-sectional area of the CL was visible within the Power Doppler sample box. After record of at least six video sequences, six images without flash artifacts and with the maximum number of coloured areas were stored from these video sequences. These images were exported in DICOM format into a computer. The CL was measured on B-mode images using a computer-assisted image analysis program (Pixelflux, version 1.0; Chameleon-Software, Leipzig, Germany); the mean value of six pictures was determined. The same software was also used to assess the total area of colour pixels within the luteal tissue. For this purpose the whole luteal structure and its blood flow area were chosen as the region of interest (ROI) and the coloured area within this ROI was calculated. The averages from four of the six images were used for further evaluation of LBF.

Blood Samples and Assays

The blood samples were obtained from the coccygeal vein by use of vacutainer vials before the applications at all days of treatments and controls. Blood samples were centrifuged immediately after collection for 15 min at 2.000 x g at room temperature; serum was stored at -20°C until assayed.

The concentrations of β C were determined by spectrophotometric analysis (Düzen Laboratory Group, TURKAK, Ankara, Turkey) and measured as described previously²⁶ after extraction using a carotene photometer (Schimadzu UV-mini 1240, Japan). Extraction efficiency was >95%. The inter-assay coefficient of variation averaged 9.7%.

Serum P_4 concentrations were measured by electrochemiluminescence immunoassay as described by Thienpont et al.²⁷ The intra- and inter-assay coefficients of variation averaged 1.4% and 2.9%, respectively.

Statistical Analysis

Statistical analyses were carried out using SPSS® Version 16.0 (SPSS Inc, Chicago, Illinois, USA). Descriptive statistics were used for determination of mean and standard error of means. Mann-Whitney U test was used for analysis of LS, LBF and blood serum β C and P_4 concentrations. Repeated Measures Define Factors test were used to determine the difference between days. A P value <0.05 was considered to be significant.

RESULTS

Animals During the Study

A total of 38 animals were synchronized and inseminated. However, only early pregnant cows were considered. Therefore, four cows in the β C- group and five cows in the β C+ group (non-pregnant animals) were withdrawn from the study. At least 15 out of 19 in the β C- group (78.94%) and 14 out of 19 in the β C+ group (73.68%), all early pregnant cows, were included in the statistical evaluation.

Beta-carotene and Progesterone Concentrations

In the β C+ group, a significant increase in serum β C concentration was determined on day 10 of gestation ($P<0.05$). However, this concentration was not maintained thereafter, especially on day 17 and 37, values were significantly lower ($P<0.01$ and $P<0.05$; Table 1). On the day of PGF₂ α treatment for induction of luteolysis, the mean serum β C concentrations of β C- and β C+ groups were determined to be 226.3 ± 105.4 μ g/dL and 229.1 ± 42.9 μ g/dL, respectively ($P>0.05$). When the mean serum β C concentration of both groups was compared to each other during early pregnancy, it was

Table 1. The mean beta-carotene concentration (μ g/dL) during early pregnancy in groups

Tablo 1. Erken gebelik döneminde grupların ortalama beta-karoten konsantrasyonları (μ g/dL)

Groups	Days					P value
	7	10	17	27	37	
β C-	280.8 \pm 50.4	289.8 \pm 42.1	226.3 \pm 43.1	287.9 \pm 48.8	272.4 \pm 49.7	n.s.
β C+	478.3 \pm 43.9 ^a	568.0 \pm 72.8 ^{b*}	390.9 \pm 88.8 ^{b***}	443.9 \pm 60.2 ^a	449.8 \pm 72.8 ^{b*}	$P^*<0.05$, $P^{**}<0.01$

n.s.: Non-significant ($P>0.05$), Values with different superscripts (a, b) in the same row are significantly different

Fig 2. Comparison of mean beta-carotene concentrations (μ g/dL) during early pregnancy between groups

Şekil 2. Erken gebelik döneminde ortalama beta-karoten konsantrasyonlarının (μ g/dL) gruplar arasında karşılaştırılması

Values with different superscripts (a, b) on the same day are different ($P^*<0.05$ and $P^{**}<0.01$)




Table 2. Comparison of mean progesterone concentrations (ng/mL) during early pregnancy between groups
Tablo 2. Erken gebelik döneminde grupların ortalama progesteron konsantrasyonları (ng/mL)

Groups	Days				
	7	10	17	27	37
βC-	2.38±0.49	6.13±1.28	9.78±1.85	13.43±3.27	14.67±3.41
βC+	4.24±2.04	8.97±2.59	7.91±1.62	14.84±1.89	13.46±3.71
P value	n.s.	n.s.	n.s.	n.s.	n.s.

n.s.: Non-significant (P>0.05)

found to be significantly higher in the βC+ group at all days investigated (Fig. 2). There was no correlation between βC concentrations, LS and LBF throughout the study.

There was no difference between groups concerning P₄ concentrations (P>0.05; Table 2).

Luteal Size

In the βC+ group, the mean LS was 4.82 cm² on day 7 and increased until day 10 (P<0.01), 17 and 37 (P<0.05). A similar increase in LS was found on day 17 and 37 of gestation in the βC- group (P<0.01; Table 3).

On the day of PGF₂α treatment, there was no significant

difference between the mean LS in the βC- and the βC+ group (4.48±0.99 cm² and 4.26±0.74 cm², respectively, P>0.05). During early pregnancy, LS was significantly higher in the βC+ group than in βC- group on day 7 (P<0.05) and 27 (P<0.01; Fig. 3).

Luteal Blood Flow

In the βC+ group, the mean LBF during pregnancy was significantly higher on day 27 and 37 than on day 7 (P<0.01). In the βC- group, it was only found higher on day 37 than on day 7 (P<0.01; Table 4).

On the day of PGF₂α injection, the mean LBF was 1.1±0.73 mm² in the βC- group and 0.62±0.27 mm² in the βC+ group,

Table 3. The mean luteal size (cm²) during early pregnancy in groups

Tablo 3. Erken gebelik döneminde gruplardaki ortalama luteal büyüklükler (cm²)

Groups	Days					P Value
	7	10	17	27	37	
βC-	4.65±1.41 ^a	4.96±0.75 ^{ab}	4.89±0.71 ^{b**}	4.19±0.99 ^{b*}	4.68±1.39 ^{b**}	P* ^{<} 0.05, P** ^{<} 0.01
βC+	4.82±1.44 ^a	5.05±0.94 ^{b**}	4.95±0.96 ^{b*}	4.73±0.69 ^{a,b}	4.80±0.88 ^{b*}	P* ^{<} 0.05, P** ^{<} 0.01

Values with different superscripts (a, b) in the same row are different

Fig 3. Comparison of mean luteal size (cm²) during early pregnancy between groups

Şekil 3. Erken gebelik döneminde ortalama luteal büyüklüklerinin (cm²) gruplar arasında karşılaştırılması

Values with different superscripts (a, b) on the same day are different (P*[<]0.05 and P**[<]0.01)




Table 4. The mean luteal blood flow (mm²) during early pregnancy in groups

Tablo 4. Erken gebelik döneminde grupların ortalama luteal kan akışkanlığı (mm²)

Groups	Days					P Value
	7	10	17	27	37	
βC-	0.61±0.32 ^a	0.84±0.42 ^{ab}	0.92±0.53 ^{a,b}	0.83±0.36 ^{a,b}	1.03±0.38 ^{b**}	P* ^{<} 0.05, P** ^{<} 0.01
βC+	0.88±0.42 ^a	0.74±0.34 ^{a,b**}	0.78±0.44 ^{a,b**}	1.21±0.65 ^{c*,**}	0.98±0.33 ^{c**}	P* ^{<} 0.05, P** ^{<} 0.01

Values with different superscripts (a, b, c) in the same row are different




Fig 4. Comparison of luteal blood flow (mm²) during early pregnancy between groups

Şekil 4. Erken gebelik döneminde luteal kan akışkanlığının (mm²) gruplar arasında karşılaştırılması

Values with different superscripts (a, b) on the same day are different ($P^* < 0.05$ and $P^{**} < 0.01$)

which was non significant ($P > 0.05$). When the mean LBF was compared between groups, it was determined to be higher in the $\beta C+$ group than in the $\beta C-$ group on day 7 only ($P < 0.05$) and 27 ($P < 0.01$; Fig. 4).

DISCUSSION

Corpus luteum is a heterogeneous tissue²⁸ consisting of steroidogenic cells (large and small luteal cells) and non-steroidogenic cells (endothelial cells, fibroblasts, pericytes and cells) originating from the bloodstream²⁹. In a complex tissue like that the various cell types must interact to ensure normal growth and development. Tissue growth depends upon growth of new blood vessels and establishment of a functional blood supply²⁸. Twenty-two percent of the mature CL consists of capillary lumina²⁹ and nearly every parenchymal cell is in contact with at least one capillary³⁰. After angiogenesis the CL becomes one of the most highly vascularised organs and receives the greatest rate of blood flow²⁸.

Local changes in blood flow within the ovary are closely related to changes in the biosynthesis of prostaglandins and steroid hormones^{31,32}. Beta-carotene is normally transported to the ovary incorporated in high density lipoprotein in the bovine³³. It is found in extremely high concentrations in the bovine CL and is responsible for the characteristic bright yellow colour of the CL³⁴. The role of βC in the bovine CL could be to support optimal steroid release³⁵. However, the relationship between serum βC and P_4 concentrations is still controversial. While many authors report no correlation³⁶⁻³⁸, others do so^{8,39}.

Many authors suppose that serum βC concentrations are affected by the nutrients taken up by cows⁴⁰⁻⁴². This study demonstrates that intramuscular βC treatments increase serum βC concentrations. In the control group, there was no significant increase in serum βC concentrations during pregnancy, and values were significantly lower than in the treatment group on days 7, 10, 27 and 37 of gestation. These findings are similar to those of others^{23,37}. On the other hand, Gossen and Hoedemaker⁴³ found that only some cows with low initial concentrations of βC ($< 200 \mu g/dL$)

showed a significant increase in serum βC concentrations after intramuscular application of βC .

However, in the present study, in the $\beta C+$ group, a significant decrease in serum βC concentrations was measured on day 17 after the last AI which resembles the findings of Ataman et al.⁴⁴ who found a significant decrease in plasma βC concentrations between day 12 and 21 of pregnancy. During early pregnancy, the up-regulation of oxytocin receptors and consecutive luteolysis is inhibited by secretion of interferon tau from the tropho-ectoderm between days 12 and 25 in cattle⁴⁵. Embryos may be transferred, and pregnancies established, as late as day 16 or 17 after estrus⁴⁶. The decrease of serum βC concentrations despite βC application might be explained by this mechanism. Eisele⁴⁷ measured high serum βC concentrations after injections of βC . Chew et al.⁴⁸ obtained high serum βC concentrations in cows after oral administration of βC . However, in the above cited publications it was reported that βC concentration significantly decreased within 10 to 14 days after the application. In our study we observed a significant decrease in serum βC concentrations 10 days (day 17 of pregnancy) after the last βC administration on day 7. On the other hand βC concentrations were maintained high for 20 days between day 17 and 37 of pregnancy, which made us suppose that the decrease prior to it might be associated with the time and period of administration.

In the present study, there were no significant differences in P_4 concentrations between groups. Some authors reported that supplementation of βC had no effect on P_4 concentration^{3,49}, however, others reported the contrary^{8,9}. It was found that P_4 concentrations in luteal tissue increased after supplementation of βC ⁵⁰. However, many fold factors such as season^{7,50} and free radicals during steroidogenesis are supposed to influence these results. In the present study, we could not assess any effect of βC treatment on serum P_4 concentrations.

Studies describing the positive effect of βC on fertility have been reported previously^{8,49,51}. In the present study, investigation of the effect of βC treatment on LS and LBF was aimed. Although a significant increase in LS during early pregnancy was observed in both groups, it was found

that the CL diameter increased more markedly in the β C+ group. On the same pregnancy days, we detected higher blood flow levels in the β C+ groups and therefore suppose that β C administration not only affects LS but in addition the LBF. It was shown previously that increasing plasma β C concentrations caused an increase in LS², and higher β C concentrations were determined in the ovaries and especially in CL by others^{2,6}, which supports our findings. The developing CL is characterized by highly active vascularisation and repeated mitoses of steroidogenic cells in parallel⁵². Vascular endothelial growth factor seems to be a major angiogenic factor responsible for vascularisation of the developing CL^{28,38}. Meanwhile, abundance of reports using colour Doppler ultrasonography for the monitoring of blood flow within the ovary and CL in cows can be found^{17,33,53}. These studies mostly revealed that there is no relation between blood flow and change in LS during the estrous cycle^{20,54}, which resembles our findings. The physiological process of CL growth and formation is probably regulated by many different factors; the mechanism is not yet completely understood⁵⁵. Recently, intraluteal mediators including cytokines and nitric oxide are thought to play some roles as pro-apoptotic and anti-apoptotic factors in the bovine CL, respectively^{56,57}. We therefore believe that β C treatment affects angiogenic factors thereby promoting vascularisation and causing the increase in LBF measured in this study.

There are several studies reporting about positive effects of β C on uterine involution and reproductive parameters⁴⁹, follicular ovulation after first postpartum follicular wave⁵⁸, embryo quality²³ and pregnancy rate³⁷. With the results we obtained in this study results we conclude that β C treatment increases serum β C concentrations, LS and LBF. The effect of β C treatment after estrus synchronisation with PGF₂ α and GnRH on fertility should be investigated in a follow up study.

ACKNOWLEDGEMENTS

The authors thank Akbel Milk and Milk Product Company Dairy Farm for allowing us to conduct this study.

REFERENCES

- Mann GE, Fray MD, Lamming GE:** Effects of time of progesterone supplementation on embryo development and interferon- τ production in the cow. *Vet J*, 171, 500-503, 2006.
- Haliloglu S, Baspinar N, Serpek B, Erdem H, Bulut Z:** Vitamin A and beta-carotene levels in plasma, corpus luteum and follicular fluid of cyclic and pregnant cattle. *Reprod Dom Anim*, 37, 96-99, 2002.
- Weng BC, Chew BP, Wong TS, Park JS, Kim HW, Lepine AJ:** β -Carotene uptake and changes in ovarian steroids and uterine proteins during the estrous cycle in the canine. *J Anim Sci*, 78, 1284-1290, 1999.
- López-Gatius F, López-Bejar M:** Reproductive performance of dairy cows with ovarian cysts after different GnRH and coloprostenol treatments. *Theriogenology*, 58, 1337-1348, 2002.
- Ahlswede L, Lotthammer KH:** Untersuchungen über eine spezifische, Vitamin A-unabhängige Wirkung des β -carotins auf die Fertilität des Rindes. 5. Mitt. Organuntersuchungen-Gewichts und Gehaltsbestimmungen. *Dtsch Tierärztl Wochenschr*, 85, 7-12, 1978.
- Arikan S, Sands H, Rodway RG, Batcheler DN:** Raman spectroscopy and imaging of β -carotene in live corpus luteum cells. *Anim Reprod Sci*, 71, 249-266, 2002.
- Aslan S, Handler J, Arbeiter K:** Gelbkörpergrösse, Progesteron, Vitamin E und β -Carotin Gehalt bei graviden Kühen. *Ankara Üniv Vet Fak Derg*, 50, 195-201, 2003.
- Aslan S, Arbeiter K, Handler J:** Frühgravidität und Embryonale bzw. Frühfetale Mortalität bei der Kuh-Gelbkörperdynamik, Progesteron, Vitamin-E, Vitamin-B12, Beta-Carotin und Folsäurekonzentrationen im peripheren blut. *Wien Tierärztl Mschr*, 85, 141-147, 1998.
- Arikan S, Rodway RG:** Effect of cyclodextrin-encapsulated β -carotene on progesterone production by bovine luteal cells. *Anim Reprod Sci*, 64, 149-160, 2000.
- McDougall S, Williamson NB, Macmilliana KL:** GnRH induces ovulation of a dominant follicle in primiparous dairy cows undergoing anovulatory follicle turnover. *Anim Reprod Sci*, 39, 205-214, 1995.
- Gümen A, Seguin B:** Ovulation rate after GnRH or PGF₂ alpha in early postpartum dairy cows. *Theriogenology*, 60, 341-348, 2003.
- Mee MO, Stevenson JS, Scoby RK:** Influence of gonadotropin-releasing hormone and timing of insemination relative to estrus on pregnancy rates of dairy cattle at first service. *J Dairy Sci*, 73, 1500-1507, 1990.
- Peters AR:** Veterinary clinical application GnRH questions of efficacy. *Anim Reprod Sci*, 88, 155-167, 2005.
- Acosta TJ, Miyamoto A:** Vascular control of ovarian function: ovulation, corpus luteum formation and regression. *Anim Reprod Sci*, 82-83, 127-140, 2004.
- Acosta TJ, Hayashi KG, Ohtani M, Miyamoto A:** Local changes in blood flow within the preovulatory follicle wall and early corpus luteum in cows. *Reprod*, 125, 759-767, 2003.
- Wiltbank MC, Dysko RC, Gallanger KP, Keyes PL:** Relationship between blood flow and steroidogenesis in the rabbit corpus luteum. *J Reprod Fert*, 84, 513-520, 1988.
- Miyamoto A, Shirasuna K, Wijayagunawardane MPB, Watabane S, Hayashi, Yamamoto D, Matsui M, Acosta TJ:** Blood flow: A key regulatory component of corpus luteum function in the cow. *Domest Anim Endocrinol*, 29, 329-339, 2005.
- Matsui M, Miyamoto A:** Evaluation of ovarian blood flow by colour Doppler ultrasound: Practical use for reproductive management in the cow. *Vet Journal*, 181, 232-240, 2009.
- Beindorf N, Nagai K, Shirasuna K, Herzog K, Hoeffmann K, Sasaki M, Bollwein H, Miyamoto A:** Vascular changes in the corpus luteum during early pregnancy in the cow. *J Reprod Dev*, 56, 263-270, 2010.
- Lüttgenau J, Ulbrich SE, Beindorf N, Honnes A, Herzog K, Bollwein H:** Plasma progesterone concentrations in the mid-luteal phase are dependent on luteal size, but independent of luteal blood flow and gene expression in lactating dairy cows. *Anim Reprod Sci*, 125, 20-29, 2011.
- Bollwein H, Baumgartner U, Stolla R:** Transrectal Doppler sonography of uterine blood flow in cows during pregnancy. *Theriogenology*, 57, 2053-2061, 2002.
- Condrad A:** Einfluss einer Supplementierung mit β -Karotin in Form einer Injektionslösung (Carofertin®) auf die Eutergesundheit von Milchkühen und die Kälbergesundheit. Zur Erhaltung des Grades einer Doktorin der Veterinärmedizin. Tierärztliche Hochschule Hannover, 2003.
- Sales JN, Dias LM, Viveiros AT, Pereira MN, Souza JC:** Embryo production and quality of Holstein heifers and cows supplemented with beta-carotene and tocopherol. *Anim Reprod Sci*, 106 (1-2): 77-89, 2008.
- Bollwein H, Prost D, Ulbrich SE, Niemann H, Honnes A:** Effects of a shortened preovulatory follicular phase on genital blood flow and endometrial hormone receptor concentrations in Holstein-Friesian cows. *Theriogenology*, 73, 242-249, 2010.
- Aslan S, Arslanbas D, Beindorf N, Bollwein H:** Effects of induction of ovulation with GnRH or hCG on follicular and luteal blood flow in Holstein-Friesian heifers. *Reprod Dom Anim*, 46, 781-786, 2011.
- Schweigert FJ, Engalbert F, Mothes R, Hurtienne A, Immig I:** Cooperative European study for the validation of a novel cow-side β -carotene assay in serum and blood. In, *Program of 13th International*

Conference on Production Disease in Farm Animals, Leipzig, Germany, p 162 (abstract), 2007.

27. Thienpont LM, Verhseghe PG, Van Brussel KA, De Leenheer AP: Efforts by industry toward standardization of serum estradiol-17 β measurements. *Clin Chem*, 44, 671-674, 1998.

28. Berisha B, Schams D: Ovarian functions in ruminants. *Domest Anim Endocrinol*, 29, 305-317, 2005.

29. Niswender GD, Juengel JL, Silva PJ, Rollyson MK, McIntush EW: Mechanisms controlling the function and life span of the corpus luteum. *Physiol Rev*, 80, 1-29, 2000.

30. Reynolds LP, Killilea SD, Redmer DA: Angiogenesis in the female reproductive system. *FASEB J*, 6, 886-892, 1992.

31. Murdoch W, Peterson T, Van Kirk E, Vincent D, Inskeep E: Interactive roles of progesterone, prostaglandins, and collagenase in the ovulatory mechanism of the ewe. *Biol Reprod*, 35, 1187-1194, 1990.

32. Acosta TJ, Yoshizawa N, Ohtani M, Miyamoto A: Local changes in blood flow within the early and midcycle corpus luteum after prostaglandin F_{2a} injection in the cow. *Biol Reprod*, 66, 651-658, 2002.

33. O'Shaughnessy PJ, Wathes DC: Bovine luteal cell activity in culture: maintenance of steroidogenesis by high density lipoprotein containing high or low beta carotene concentrations. *Anim Reprod Sci*, 17, 165-176, 1988.

34. O'Fallon JV, Chew BP: The subcellular distribution of β -carotene in bovine corpus luteum. *Proc Soc Exp Biol Med*, 177, 406-411, 1984.

35. Arikan S, Rodway RG: The effect of beta-carotene either in organic solvents or as HDL on progesterone production by bovine luteal cells. *J Reprod Fertil*, 19, 64, 1997.

36. Folman Y, Ascarelli I, Herz Z, Rosenberg M, Davidson M, Haleiv A: Fertility of dairy heifers given a commercial diet free of β -carotene. *Br J Nutr*, 41, 353-359, 1979.

37. Aréchiga CH, Staples CR, McDowell LR, Hansen PJ: Effects of timed insemination and supplemental β -Carotene on reproduction and milk yield of dairy cows under heat stress. *J Dairy Sci*, 81, 390-402, 1998.

38. Ferreira-Dias GM, Skarzynski DJ: Some aspects of regulation of luteal function and luteolysis in equine corpora lutea. *Pferdeheilkunde*, 24, 10-14, 2008.

39. Lothammer KH: Importance and role of beta-carotene for bovine fertility. In, Importance of Beta-carotene for Bovine Fertility Roche Symposium, Roche, London, UK, pp. 5-44, 1978.

40. Bindas EM, Gwazdauskas FC, Aiello RJ, Herbein JH, McGilliard ML, Polan CE: Reproductive and metabolic characteristics of dairy cattle supplemented with β -carotene. *J Dairy Sci*, 67, 1249-1255, 1984.

41. Wang JY, Owen FG, Larson LL: Effect of carotene supplementation on reproductive performance lactating Holstein cows. *J Dairy Sci*, 71, 181-186, 1988.

42. Kolb E, Seehawer UJ: Die Bedeutung der Carotine und des Vitamin A für die Fortpflanzung bei Rindern, Pferden und Schweinen - eine Übersicht. *Prakt Tierarzt*, 78, 783-789, 1997.

43. Gossen N, Hoedemaker M: Effect of beta-carotene serum concentration

on the reproductive performance in dairy cows. *Berl Munch Tierarztl Wochenschr*, 118, 326-333, 2005.

44. Ataman MB, Erdem H, Bülbül B, Haliloğlu S, Çınar M, Aköz M: Plasma β -carotene, vitamin A and vitamin C levels in cyclic and pregnant cows. *Kafkas Univ Vet Fak Derg*, 16 (4): 579-584, 2010.

45. Farin CE, Imakawa K, Hansen TR, McDonnell JJ, Murphy CN, Farin PW, Roberts RM: Expression of trophoblastic interferon genes in sheep and cattle. *Biol Reprod*, 43, 210-218, 1990.

46. Betteridge KJ, Eaglesome MD, Randall GCB, Mitchell D: Collection, description and transfer of embryos from cattle 10-16 days after oestrus. *J Reprod Fertil*, 59, 205-216, 1980.

47. Eisele W: Beta-Carotin und Vitamin A in Blut und Milch des Rindes nach parenteraler und oraler Applikation von Beta-Carotin. Tierärztliche Fakultät PhD Thesis, München Univ (Germany), 1987.

48. Chew BP, Wong TS, Michal JJ: Uptake of orally administered β -carotene by blood plasma, leukocytes, and lipoproteins in calves. *J Anim Sci*, 71, 730-739, 1993.

49. Kaewlamun W, Okuyi M, Humblot P, Techakumphu M, Ponter AA: Does supplementing dairy cows with β -carotene during the dry period affect postpartum ovarian activity, progesterone, and cervical and uterine involution? *Theriogenology*, 75, 1029-1038, 2011.

50. Graves-Hoagland RL, Hoagland TA, Woody CO: Effect of β -carotene and vitamin A on progesterone production by bovine luteal cells. *J Dairy Sci*, 71, 1058-1062, 1988.

51. Iwanska S, Lewicki C, Rybicka M: The effect of beta carotene supplementation on the beta carotene and vitamin A levels of blood plasma and some fertility indices of dairy cows. *Arch Tierernähr*, 35, 563-570, 1985.

52. Reynolds LP, Grazul-Bilska AT, Redmer DA: Angiogenesis in the corpus luteum. *Endocrine*, 12, 1-9, 2000.

53. Ginther OJ, Silva LA, Araujo RR, Beg MA: Temporal associations among pulses of 13,14-dihydro-15-keto-PGF₂ α , luteal blood flow, and luteolysis in cattle. *Biol Reprod*, 76, 506-513, 2007.

54. Herzog K, Bollwein H: Application of Doppler ultrasonography in cattle reproduction. *Reprod Dom Anim*, 42 (Suppl 2): 51-58, 2007.

55. Skarzynski DJ, Ferreira-Dias G, Okuda K: Regulation of luteal function and corpus luteum regression in cows: Hormonal control, immune mechanisms and intercellular communication. *Reprod Dom Anim*, 43 (Suppl 2): 57-65, 2008.

56. Petroff MG, Petroff BK, Pate JL: Mechanisms of cytokine-induced death of cultured bovine luteal cells. *Reproduction*, 121, 753-760, 2001.

57. Skarzynski DJ, Jaroszewski JJ, Okuda K: Role of tumor necrosis factor- α and nitric oxide in luteolysis in cattle. *Domest Anim Endocrinol*, 29, 340-346, 2005.

58. Kawashima C, Kida K, Schweigert FJ, Miyamoto A: Relationship between plasma β -Carotene concentrations during the peripartum period and ovulation in the first follicular wave postpartum in dairy cows. *Anim Reprod Sci*, 111, 105-111, 2009.