



## RESEARCH ARTICLE

### The mismatched isolation of *Brucella* strains from nomic hosts

Özgür Çelebi<sup>1\*</sup>, Fatih Büyük<sup>1</sup>, Ketı Sidamonidze<sup>2</sup>, Eka Zhgenti<sup>2</sup>, Doğan Akça<sup>3</sup>, Mitat Şahin<sup>1</sup>

<sup>1</sup>Department of Microbiology, Faculty of Veterinary Medicine, Kafkas University, Kars, 36100, Turkey, <sup>2</sup>Department of Virology, Molecular Biology and Genome Research, R.G. Lugar Center for Public Health Research, National Center for Disease Control and Public Health, Tbilisi, 0177, Georgia, <sup>3</sup>Kars Health School, University of Kafkas, Kars, 36100, Turkey  
Geliş: 17.10.2016, Kabul: 05.12.2016  
\*ozgurcelebi36@hotmail.com

#### Öz

**Çelebi Ö, Büyük F, Sidamonidze K, Zhgenti E, Akça D, Şahin M.** Primer konaklardan izole edilen uyumsuz *Brucella* türleri.

#### Abstract

**Celebi O, Buyuk F, Sidamonidze K, Zhgenti E, Akca D, Sahin M.** The mismatched isolation of *Brucella* strains from nomic hosts.

Eurasian J Vet Sci, 2017, 33, 1, 40-45

DOI: 10.15312/EurasianJvetSci.2016.134

**Amaç:** Bu çalışmada Kafkas Üniversitesine ait *Brucella* suş koleksiyonunun bir bölümü çeşitli yöntemler ile değerlendirilmiştir.

**Gereç ve Yöntem:** Suşların orjini abort hikayesi olan sığır ve koyunlar oluşturdu. Otuz adet *Brucella* suşu (16'sı koyun ve 14'ü sığır orjinli) rastgele seçildi ve bu çalışmaya dahil edildi. Standardize fenotipik karakterizasyona ilaveten suşların moleküler tiplendirmesi için cins spesifik Real Time PCR ve tür spesifik Bruce-Ladder PCR uygulandı.

**Bulgular:** Onaltı koyun örneğinin 15'i *B. melitensis* ve 14 sığır örneğinin 13'ü *B. abortus* olarak tanımlandı. İlginçtir ki, koyun aborte fütustan izole edilen bir suş *B. abortus* ve sığırdan izole edilen bir suş *B. melitensis* olarak tanımlandı. Bununla birlikte, koyun fütusundan elde edilen bir suş bütün fenotipik özellikleri yönünden *Brucella* cinsine benzer karakterde olmasına rağmen genotipik olarak (ne Real Time PCR ne de Bruce-Ladder PCR) *Brucella* spp. olarak doğrulanamadı ve ileride karakterize edilmek üzere *Ochrobactrum* spp. şüpheli olarak saklandı.

**Öneri:** Bu çalışma belli konaklar arasında çapraz *Brucella* enfeksiyonunun sıradışı olarak nitelendirilebileceğini fakat zamanla bunun olağan olabileceğini göstermektedir. Bu şekildeki çalışmalar, *Brucella* cinsinin patojenitesini, konak spesifitesini ve gelişimini daha iyi anlamaya yardımcı olacaktır.

**Anahtar kelimeler:** *Brucella*, saha suşlar, sığır, koyun, moleküler karakterizasyon

**Aim:** A small part of field *Brucella* strain collection of Kafkas University was evaluated by different methods in this study.

**Materials and Methods:** The strains were originated from cattle and sheep which had a history of abortion. Thirty field *Brucella* strains (16 were originated from sheep and 14 were from cattle) were randomly selected and included in the study. In addition of standardized phenotypical characterization, genus specific real-time PCR and species specific Bruce-Ladder PCR were conducted for molecular typing.

**Results:** Out of 16 sheep and 14 cattle samples 15 and 13 were identified as *B. melitensis* and *B. abortus*, respectively. Interestingly one strain was characterized as *B. abortus* isolated from aborted sheep foetus and the other as *B. melitensis* from aborted cattle milk sample. Furthermore one strain from aborted sheep foetus was unable to be confirmed as *Brucella* spp. by genotypically (neither by real-time PCR nor Bruce-Ladder PCR) though all phenotypic features were similar to *Brucella* genus and kept under strict conditions for future characterization on suspicion of *Ochrobactrum* spp.

**Conclusions:** These report alleges that *Brucella* cross-infection among certain hosts ceased to be extraordinary and gained an usual face over time. These kind of efforts would help in further understanding the evolution, host specificity and pathogenicity of the genus *Brucella*.

**Keywords:** *Brucella*, field strains, cattle, sheep, molecular characterization



## Introduction

*Brucella* spp. are the causative agents of brucellosis, an infectious disease that affects various species of animals and can be transmitted to humans through direct contact with infected animals, indirectly by the ingestion of raw milk products, and during the handling of strains or infected material in the laboratory (Otlu et al 2008, Aras et al 2009). *Brucellosis* is endemic in Turkey and causes severe economic losses in livestock. Studies in various parts of the country indicate that the disease is widespread among cattle and sheep populations (İyisan et al 2000, İça et al 2014). *Brucellosis* in farm animals have been reported in East Anatolian Region in where Kars is the most important city with the huge animal population and substantial amount cases of given disease (Ünver et al 2006, Şahin et al 2008, Çelebi et al 2011, Büyük et al 2011).

Currently the *Brucella* genus consists of 12 species with validly published names, based on host preferences, phenotypic differences and pathogenesis (Godfroid et al 2011, Whatmore et al 2014, Scholz et al 2016). While a broad host range generally exists for *Brucella* species, *Brucella* infection follows a very strict, host-related hierarchy of pathogenicity (Adam 2002). It is known that the principal species of *Brucella* in cattle and sheep are *B. abortus* and *B. melitensis*, respectively. Some individual biotypes has limited scale and cross-infection among host species is very rare in natural condition. Inappropriate management may allow the disease to be transferred to a heterologous host such as cattle and sheep. Unexpected isolation of related strains from both hosts are likely and may also causes occasional infections in animals resulting in abortions (Ocholi et al 2005, Büyükcan-gaz and Şen 2007, Büyük et al. 2011, Erdenliğ Gürbilek et al 2014).

In this study, a small part of field *Brucella* strain collection of Kafkas University was evaluated by genus specific Real-Time PCR and species specific Bruce-Ladder PCR in addition of standardized phenotypical characterization.

## Materials and Methods

### Bacterial isolation and identification

In this study, field *Brucella* strains were evaluated by bacteriological and molecular methods. For this purpose a small part of culture collection of Microbiology Department of Veterinary Faculty, Kafkas University, Kars (Turkey) was used. The collection was developed from the samples taken from the cattle and sheep which had a history of abortion and human who were in a febrile disease in Kars region. The collection totally includes 259 *Brucella* strains that were composed 115 sheep, 139 cattle and 5 human isolates. Thirty field strains of *Brucella* sp. were randomly selected and included in the study as originated 16 isolates from sheep (all were from

aborted foetus) and 14 from cattle (4 were from aborted foetus, 5 from milk and 5 from vaginal secret).

Bacterial isolation was made from the samples such as aborted foetus, milk and vaginal secret were duly taken and handed meticulously in accordance with the report of Büyük and Şahin (2011). In brief, milk and vaginal secret samples were first enriched in Farrel Broth for 5-7 days. Then the content was plated on *Brucella* Selective Agar (Oxoid, UK) plates containing *Brucella* selective supplement (Oxoid, UK). The fetal tissue samples were directly plated on selective agar plates. All plates were incubated at atmospheric condition with and without 5-10% CO<sub>2</sub> at 37°C for 5-7 days.

For the identification of the genus and/or species of *Brucella*, cultural and biochemical tests, including colony and microscopic morphology, catalase, oxidase, urease tests, H<sub>2</sub>S production, Tbilisi phage susceptibility and CO<sub>2</sub> requirement were performed (Alton et al 1988).

### Molecular typing

For molecular typing, genus-specific real-time PCR and species-specific BruceLadder-PCR were conducted. Genomic DNA was extracted from pure cultures of strains through the commercial extraction kit (Qiagen, Germany) in line with the manufacturer's recommendations.

Real-time was carried out with Freeze-Dried Reagent Kit specific for *Brucella* genus. The kit is specially adapted for amplification in glass capillaries using the Idaho Technoogy's Ruggedized Advanced Pathogen Identification Device (R.A.P.I.D.) Instruments and TaqMan probes. The kit includes negative control (NC), unknown sample assays, positive control (PC) vials, 2X Reconstitution Buffer and Reagent Grade Water. Each reagent vial contains enough master mix for two reactions. 20 µL of 2X Reconstitution Buffer was added on to the NC, unknown and PC reagent vials, respectively. Twenty µL of Reagent Grade Water was added on to the NC and PC reagent vials. Twenty µL of sample (template DNA) was added on to the unknown reagent vial as well. After vortex and centrifugation of all tubes, 19 µL of the hydrated mixture was transferred into a capillary in duplicate. Once the samples were prepared the protocols R.A.P.I.D. 7200 was followed in LightCycler 2.0 Instrument.

For confirmation of *Brucella* species Bruce-Ladder PCR was carried out by modified from Garcia-Yoldi et al. (2006) (Table 1).

*Brucella* reference strains (*B. abortus* biotype 1 (544) and *B. melitensis* biotype 1 (16M)) and negative control (*E. coli* OP50) were used in both PCR. The amplified products were resolved by electrophoresis using a 1.5% agarose gel and followed by staining with ethidium bromide.





## Results

The reaction content and *Brucella* species-specific Bruce-Ladder PCR conditions and sources and results of samples tested by conventional and molecular methods were displayed in Table 1 and Table 2. Results of *Brucella* genus-specific real-time PCR and *Brucella* species-specific PCR (Bruce-Ladder-PCR) in Figure 1A ad 1B. All 30 field strains were found as *Brucella* spp. by cultural and biochemical tests. Out of 16 sheep samples and 14 cattle originated samples 15 and 13 were identified as *B. melitensis* and *B. abortus*, respectively. Interestingly one strain (#3) was characterized as *B. abortus* isolated from aborted sheep foetus and one (#19) as *B. melitensis* from aborted cattle milk sample

All strains previously identified as *Brucella* spp. by conventionally were future confirmed as *Brucella* spp. with the genus-specific real-time PCR. The expression level of Target 1 region related to the all certain strains of *Brucella* genus was measured using an average crossing point (Cp) of each sample that was run in duplicate. And Cp values were found with a range of 13,69 and 14,44 when compared the Cp value of PC as 21,59. The NC and one strain (#28) didn't give any significant Cp values.




Figure 1 (A-B): Results of *Brucella* genus and species-specific PCR (Bruce-Ladder PCR). (A) Results of *Brucella* genus-specific real-time PCR, blue and green curves; field *B. abortus* (#22), red and black curves; field *B. melitensis* (#173), pink and dark green; negative control (NC), dark blue and grey; uncharacterized strain (#312); pale red and purple; positive control (PC). (B) Results of *Brucella* species-specific PCR (Bruce-Ladder-PCR), lane 1; positive control (*B. melitensis* 16M), lane 2; positive control (*B. abortus* 544), lane 3; field *B. melitensis* isolate, lane 4; field *B. abortus* isolate, lane5; molecular weight stand (100 bp DNA ladder, New England BoiLab).

Fifteen isolates of *B. abortus* and 13 of *B. melitensis* presented a common Bruce-Ladder PCR profile expected for field strains, excluding the vaccine strains, with the several band sizes of 1682, 794, 587, 450 and 152bp, except the size of 1071bp that is specific for only *B. melitensis* (Figure 1B). Interestingly one strain (#19) of cattle and one strain (#3) of sheep presented unusual Bruce-Ladder PCR band profiles specific for *B. melitensis* and *B. abortus*, respectively.

Furthermore one strain (#28) from aborted sheep foetus was unable to be identified as *Brucella* spp. by genotypically (neither by real-time PCR nor Bruce-Ladder PCR) though all phenotypic features are very similar to *Brucella* genus. This strain was suspected to be one of *Ochrobactrum* species and kept under strict conditions for future characterization.

## Discussion

In comparison with the broad host range of *Brucella* species exist, the infection has very strict host specificity. Thus, goats are the natural hosts of *B. melitensis* and sheep are preferred hosts of the pathogen, and cattle are the natural hosts of *B. abortus*. In some cases, the certain *Brucella* species cross the border from its natural host, transmit to the others and lead to exceptional abortion in unfavorable hosts (Ocholi et al 2005). Contrary to this, the isolation of *B. melitensis* from cattle has been reported constantly during a decade (Refai 2002) and it substituted for *B. abortus* in cattle in South Europe (Godfroid and Kasbohrer 2002). Similar habit was reported for *B. abortus* as a causative agent of sheep brucellosis in both natural and experimental conditions in some regions (Shaw 1976, Zowghi and Ebadi 1988, Ocholi et al 2005, Sha-reef 2006).

Though it was scarce of cross-infection in interspecies it is possible due to the common use of shelters and pastures by different animal like in the reported study. This kind of mismatched isolation was first reported virtually by Büyük-cangaz and Şen (2007) in cattle in Turkey and the causative agent was typed as *B. melitensis* biotip 3. A different study

Table 1. *Brucella* species-specific BruceLadder-PCR conditions.

PCR reagents		PCR reaction thermal conditions and cycles			
Reagent	Vol.(µL)	Cycle element	Temp (°C)	Time (sec)	# Cycles
PCR buffer (x10)	2.5	Initial denaturation	95	35	1
dNTP mix (2.5 mM)	2.5	Denaturation	95	35	35
MgCl <sub>2</sub> (50 mM)	1.5	Annealing	63	45	
Bruceladder Primer mix (10 pmol)	2	Elongation	72	180	
Taq DNA polymerase (5u/ul)	0.2	Final elongation	72	360	1
Water	15.3				
Templete DNA (10 ng/ul)	1				



Table 2. Sources and results of samples tested by conventionally and molecular methods.

Sample number	Origin	Sample type	Time originally isolated	Phenotypic characterization						Genotyping c haracterization			Final result
				CO2 req.	H2S	Catalase	Oxidase	Urease	Tbilisi Phage	Bruce-			
										Real-time PCR	Ladder PCR		
#1	Sheep	Foetus	2005	-	--	+	+	+	-	+	+	<i>B. melitensis</i>	
#2	Cattle	Foetus	2005	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#3	Sheep	Foetus	2005	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#4	Sheep	Foetus	2006	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#5	Sheep	Foetus	2007	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#6	Cattle	Foetus	2007	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#7	Sheep	Foetus	2007	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#8	Sheep	Foetus	2008	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#9	Sheep	Foetus	2009	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#10	Sheep	Foetus	2009	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#11	Sheep	Foetus	2009	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#12	Cattle	Milk	2009	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#13	Cattle	Milk	2009	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#14	Cattle	V. secret	2009	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#15	Cattle	V. secret	2009	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#16	Cattle	Milk	2008	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#17	Cattle	V. secret	2008	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#18	Cattle	Milk	2010	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#19	Cattle	Milk	2010	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#20	Sheep	Foetus	2010	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#21	Cattle	V. secret	2010	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#22	Cattle	V. secret	2010	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#23	Sheep	Foetus	2011	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#24	Cattle	Foetus	2011	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#25	Sheep	Foetus	2011	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#26	Sheep	Foetus	2012	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#27	Cattle	Foetus	2013	+	+	+	+	+	+	+	+	<i>B. abortus</i>	
#28	Sheep	Foetus	2013	-	-	+	+	+	-	-	-	<i>Uncharacterized</i>	
#29	Sheep	Foetus	2013	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	
#30	Sheep	Foetus	2014	-	-	+	+	+	-	+	+	<i>B. melitensis</i>	

DİP: Deneme intraperitoneal, KİP: Kontrol intraperitoneal, DİM: Deneme immersiyo, KİM: Kontrol immersiyo

was from Kars region followed this with a report of case more than half *Brucella* isolates originated from cattle samples were representing as if PCR profile that was very similar to *B. melitensis* vaccine strain-Rev 1 (Unver et al 2006). Another exceptional case was reported by Büyük and Şahin (2011) for Kars strains of which one strain was isolated from aborted cattle foetus and identified as *B. melitensis* subsequ-

ently. This kind of reports are leaving away especially for *B. melitensis* as unusual causative agent of cattle brucellosis in Turkey (Sarisayin et al 1969, Erdenliğ Gürbilek et al 2014). Conversely sheep cases arising from *B. abortus* are more rarely in Turkey (Erdenliğ Gürbilek et al 2014) even it was reported many countries heretofore (Zowghi and Ebadi 1988, Ocholi et al 2005, Shareef 2006).







In this study one strain was characterized as *B. abortus* isolated from aborted sheep fetus and one as *B. melitensis* from aborted cattle milk sample. The results are close similarity and the explanation of this kind of unusual case can illustrated as in the studies reported before; having common habitats of animals, lack of an effective vaccine for cross-protection and fully optimized precaution method of disease and limited knowledge on the epidemiology in these host species (Pishva and Salehi 2008, Alvarez et al 2011, Büyük and Şahin 2011). But the important that it should not be ignored of reporting of *Brucella* cross-infection among certain hosts ceased to be extraordinary and gained an usual face over time.

An other interesting result is an unusual strain (#28) isolation from aborted sheep fetus that was unable to be confirmed as *Brucella* spp. by genotypically though all phenotypic features (aerobic, motile, oxidase and urease positive, Gram-negative rods) are very similar to *Brucella* genus. This strain was suspected to be one of *Ochrobactrum* species based on its features. This kind of close relationship has been reported between *B. melitensis* and *Ochrobactrum* spp. and has also led to misidentification (Elsaghir and James 2003). Besides this phenotypic features *O. anthropi* is one of the closest *Brucella* relatives based on DNA, rRNA, and protein analyses (Yanagi and Yamasato 1993, Velasco et al 1998, Cloeckaert et al 1999). It is known that some of *Ochrobactrum* strains can cause severe systemic infections in healthy human opportunistically and mimics *Brucella* infection (Kettaneh et al 2003, Ozdemir et al 2006). Despite all, *Ochrobactrum* species are unable to establish a chronic infection and have never associated with abortive animal infections. But it is still kept under strict conditions for future characterization and on probation as an unusual inductive agent for abortive animal infection.

## Conclusions

In conclusion, owing to the ability of *Brucellae* to adapt to different conditions that leads to changes in epidemiologic features of *Brucella* species it would be advantageous to reveal strain diversity and illuminate complexity of interactions between the organism and animals. No doubt that we will continue to expand our strain pool and to full characterization of our remaining *Brucella* strains and thus will be able to make out these questions. These laborious approaches should help in further understanding the evolution, host specificity and pathogenicity of the genus *Brucella*. Thus the preliminary results would be beneficial and will attract attention of authorized persons to improve control measurements and even encourage them to contribute of production of novel polyvalent vaccine that is able to protect both hosts.

## Acknowledgments

We would like to thank the Kafkas Universtiy, Department of Microbiology persons who participated in the study for their supports.

## References

- Adams LG, 2002. The pathology of brucellosis reflects the outcome of the battle between the host genome and the *Brucella* genome. *Vet Microbiol*, 90, 553-561.
- Alton GG, Jones LM, Angus RD, Verger JM, 1988. Techniques for the brucellosis laboratory. INRA (Institut National de la Recherche Agronomique), Paris, France.
- Alvarez J, Saez JL, Garcia N, Serrat C, Perez-Sancho M, Gonzalez S, Ortega MJ, Gou J, Carbajo L, Garrido F, Goyache J, Dominguez L, 2011. Management of an outbreak of brucellosis due to *B. melitensis* in dairy cattle in Spain. *Res Vet Sci*, 90, 208-211.
- Aras Z, Ateş M, Uçan US, 2009. *Brucella* suşlarının identifikasyonu ve biyotiplendirilmesi. *Eurasian J Vet Sci*, 25, 51-59.
- Büyükcangaz E, Şen A, 2007. The first isolation of *Brucella melitensis* from bovine aborted fetus in Turkey. *JBES*, 1, 139-142.
- Büyük F, Şahin M, 2011. Investigation of *Brucella* species from various samples of aborted cattle in Kars Province (Turkey) by cultural and molecular methods and epidemiological analysis of cases. *Kafkas Üniv Vet Fak Derg*, 17, 809-816.
- Büyük F, Çelebi Ö, Şahin Ö, Ünver A, Tazegül E, 2011. *Brucella* and *Campylobacter* mixed infection in two different sheep and goat herds. *Kafkas Üniv Vet Fak Derg*, 17, 177-180.
- Cloeckaert A, Tibor A, Zygmunt MS, 1999. *Brucella* outer membrane lipoproteins share antigenic determinants with bacteria of the family Rhizobiaceae. *Clin Diagn Lab Immunol*, 6, 627-629.
- Çelebi Ö, Otlı S, 2011. Bacteriological and Molecular Description of *Brucella* Species Isolated from Milk and Vaginal Swab Samples of Aborted Cattle in Kars Region. *Kafkas Üniv Vet Fak Derg*, 17, 53-58.
- Elsaghir AA, James EA, 2003. Misidentification of *Brucella melitensis* as *Ochrobactrum anthropi* by API 20NE. *JMM*, 52, 441-442.
- Erdenliğ Gürbilek S, Baklan EA, Aksoy HY, 2014. Identification and phage susceptibilities of *Brucella* strains in Turkey between the years 2007 and 2008. *Harran Üniv Vet Fak Derg*, 3, 67-72.
- Garcia-Yoldi D, Marin CM, de Miguel MJ, Munoz PM, Vizmanos JL, Lopez-Goni I, 2006. Multiplex PCR assay for the identification and differentiation of all *Brucella* species and the vaccine strains *Brucella abortus* S19 and RB51. *Clin Chem*, 52, 779-781.
- Godfroid J, Käsbohrer A, 2002. Brucellosis in the European Union and Norway at the turn of the twenty-first century. *Vet Microbiol*, 90, 135-145.



- Godfroid HC, Scholz T, Barbier C, Nicolas C, Wattiau P, Fretin D, Whatmore AM, Cloeckaert A, Blasco JM, Moriyon I, Sagerman C, Muma JB, Al Dahouk S, Neubauer H, Letesson JJ, 2011. Brucellosis at the animal/ecosystem/human interface at the beginning of the 21st century. *Prev Vet Med*, 102, 118-131.
- Iyisan AS, Akmaz O, Duzgun SG, Ersoy Y, Eskiizmirli S, Güler L, Gündüz K, Işık N, İçyerioğlu AK, Kalender H, Karaman Z, Küçükayan U, Özcan C, Seyitoğlu S, Tuna D, Tunca T, Üstünakın K, Yurtalan, S. 2000. Türkiye’de sığır ve koyunlarda Brucellosis’in seroepidemiolojisi. *Pendik Hay Hast M Araş Enst Derg*, 31, 21-75.
- İça T, Aydın F, Gümüşsoy KS, Perçin D, Sümerkan AB, Ocak F, Abay S, Doğan HO, Fındık A, Çiftçi A, 2012. Conventional and molecular biotyping of *Brucella* strains isolated from cattle, sheep and human. *Ankara Üniv Vet Fak Derg*, 59, 259-264.
- Kettaneh A, Weill FX, Poilane I, Fain O, Thomas M, Herrmann JL, Hocqueloux L, 2003. Septic shock caused by *Ochrobactrum anthropi* in an otherwise healthy host. *JCM*, 41, 1339-1341.
- Ocholi RA, Kwaga JK, Ajogi I, Bale JO, 2005. Abortion due to *Brucella abortus* in sheep in Nigeria. *Rev Sci Tech*, 24, 973-979.
- Otlı S, Şahin M, Atabay Hİ, Ünver A, 2008. Serological investigations of brucellosis in cattle, farmers and veterinarians in the Kars District of Turkey. *Acta Vet Brno*, 77, 117-121.
- Özdemir D, Soyupacı Z, Şahin I, Bicik Z, Şencan I, 2006. *Ochrobactrum anthropi* endocarditis and septic shock in a patient with no prosthetic valve or rheumatic heart disease: case report and review of the literature. *JJID*, 59, 264-265.
- Pishva E, Salehi M, 2008. First report of isolation of *Brucella melitensis* vaccine Rev.1 and a source of cattle infection in Iran. *J Sci I R Iran*, 19, 19-23.
- Refai M, 2002. Incidence and control of brucellosis in the Near East region. *Vet Microbiol*, 90, 81-110.
- Sarısayın F, Eroğlu M, Nadas UG, 1969. Yurdumuzda izole edilen Brusella suşlarının tür ve biyotiplerini tayini ile dağılışı durumu üzerine bir çalışma. *Pendik Hay Hast M Araş Enst Derg*, 1, 24-35.
- Scholz HC, Revilla-Fernandez S, Al-Dahouk S, Hammerl JA, Zygmunt MS, Cloeckaert A, Koylass M, Whatmore AM, Blom J, Vergnaud G, Witte A, Aistleitner K, Hofer E, 2016. *Brucella vulpis* sp. nov., isolated from mandibular lymph nodes of red foxes (*Vulpes vulpes*). *Int J Syst Evol Microbiol*, 66, 2090-2098.
- Shareef JM, 2006. Isolation of *Brucella abortus* from aborted lambs in the governorate of Sulaimania/Said-Sadic District. *Iraqi J Vet Sci*, 20, 19-23.
- Shaw WB, 1976. *Brucella abortus* infection in sheep. II. Experimental infection of ewes. *Br Vet J*, 132, 143.
- Şahin M, Genç O, Ünver A, Otlı S, 2008. Investigation of bovine brucellosis in the Northeastern Turkey. *Trop Anim Health Prod*, 40, 281-286.
- Ünver A, Erdoğan HM, Atabay Hİ, Şahin M, Çelebi Ö, 2006. Isolation, identification, and molecular characterization of *Brucella melitensis* from aborted sheep fetuses in Kars, Turkey. *Rev Med Vet*, 157, 42-46.
- Ünver A, Erdoğan HM, Atabay Hİ, Şahin M, Güneş V, Çitil M, Gökçe Hİ, 2006. Genotyping of *Brucella* spp. isolated from aborted cattle fetuses by RAPD-PCR. *Kafkas Üniv Vet Fak Derg*, 12, 121-127.
- Velasco J, Romero C, Lopez-Goni I, Leiva J, Diaz R, Moriydn I, 1998. Evaluation of the relatedness of *Brucella* spp. and *Ochrobactrum anthropi* and description of *Ochrobactrum intermedium* sp. nov., a new species with a closer relationship to *Brucella* spp. *Int J Syst Bact*, 48, 759-768.
- Whatmore AM, Davison N, Cloeckaert A, Al Dahouk S, Zygmunt MS, Brew SD, Perrett LL, Koylass MS, Vergnaud G, Quance C, Scholz HC, Dick EJ Jr, Hubbard G, Schlabritz-Loutsevitch NE, 2014. *Brucella papionis* sp. nov., isolated from baboons (*Papio* spp.). *Int J Syst Evol Microbiol*, 64, 4120-4128.
- Yanagi M, Yamasato K, 1993. Phylogenetic analysis of the family Rhizobiaceae and related bacteria by sequencing of 16S rRNA gene using PCR and DNA sequencer. *FEMS Microbiol Lett*, 107, 115-120.
- Zowghi E, Ebadi A, 1988. Abortion due to *Brucella abortus* in sheep in Iran. *Rev Sci Tech*, 7, 379-382.

