Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat.
Volume 66, Number 1, Pages 85–91 (2017)
DOI: 10.1501/Commua1_0000000777
ISSN 1303–5991

INITIAL COEFFICIENTS FOR A SUBCLASS OF BI-UNIVALENT FUNCTIONS DEFINED BY SALAGEAN DIFFERENTIAL OPERATOR

MURAT ÇAĞLAR AND ERHAN DENIZ

Abstract. In this paper, we investigate a new subclass $\Sigma^n(\tau,\gamma,\varphi)$ of analytic and bi-univalent functions in the open unit disk $\mathcal{U}$ defined by Salagean differential operator. For functions belonging to this class, we obtain estimates on the first two Taylor-Maclaurin coefficient $|a_2|$ and $|a_3|$.

1. INTRODUCTION

Let $\mathcal{A}$ denote the class of functions f of the form

$$f(z) = z + \sum_{k=2}^{\infty} a_k z^k \tag{1.1}$$

which are analytic in the open unit disk $\mathcal{U} = \{z : |z| < 1\}$. We also denote by $\mathcal{S}$ the class of all functions in $\mathcal{A}$ which are univalent in $\mathcal{U}$.

Salagean [18] introduced the following differential operator for $f(z) \in \mathcal{A}$ which is called the Salagean differential operator:

$$\begin{aligned} D^0 f(z) &= f(z) \\ D^1 f(z) &= Df(z) = zf'(z) \\ D^n f(z) &= D(D^{n-1} f(z)) \quad (n \in \mathbb{N} = 1, 2, 3, ...). \end{aligned}$$

We note that,

$$D^n f(z) = z + \sum_{k=2}^{\infty} k^n a_k z^k \qquad (n \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}). \tag{1.2}$$

It is well known that every $f \in \mathcal{S}$ has an inverse function f^{-1} satisfying

$$f^{-1}(f(z)) = z \quad (z \in \mathcal{U})$$

Received by the editors: May 04, 2016, Accepted: Aug. 25, 2016.
2010 *Mathematics Subject Classification.* Primary 30C45, 30C50; Secondary 30C80.
Key words and phrases. Analytic functions, Univalent functions, Bi-univalent functions, Subordination, Salagean differential operator.

©2017 Ankara University

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f);\ r_0(f) \geq \frac{1}{4}\right)$$

where

$$f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + ...$$

A function $f \in \mathcal{A}$ is said to be bi-univalent in $\mathcal{U}$ if both $f(z)$ and $f^{-1}(z)$ are univalent in $\mathcal{U}$. Let Σ denote the class of bi-univalent functions in $\mathcal{U}$ given by (1.1). Lewin [13] introduced the bi-univalent function class and showed that $|a_2| < 1.51$. Subsequently, Brannan and Clunie [2] conjectured that $|a_2| \leq \sqrt{2}$. Netanyahu [15], otherwise, showed that $\max_{f\in\Sigma}|a_2| = \frac{4}{3}$. The coefficient estimate problem for each of the following Taylor Maclaurin coefficients: $|a_n|$ $(n \in \mathbb{N}\setminus\{1,2\}\,;\ \mathbb{N} = \{1,2,3,...\})$ is still an open problem. Recently, several researchers such as ([1]-[7], [9]-[16], [17], [19]-[24]) obtained the coefficients $|a_2|$, $|a_3|$ of bi-univalent functions for the various subclasses of the function class Σ . Motivating with their work, we introduce a new subclass of the function class Σ and find estimates on the coefficients $|a_2|$ and $|a_3|$ for functions in these new subclass of the function class Σ employing the techniques used earlier by Srivastava et al. [19] and Frasin and Aouf [9].

Let φ be an analytic and univalent function with positive real part in $\mathcal{U}$, $\varphi(0) = 1$, $\varphi'(0) > 0$ and φ maps the unit disk $\mathcal{U}$ onto a region starlike with respect to 1 and symmetric with respect to the real axis. The Taylor's series expansion of such function is

$$\varphi(z) = 1 + B_1z + B_2z^2 + B_3z^3 + ..., \tag{1.3}$$

where all coefficients are real and $B_1 > 0$. Throughout this paper we assume that the function φ satisfies the above conditions unless otherwise stated.

Definition 1.1. *A function $f \in \Sigma$ given by (1.1) is said to be in the class $\Sigma^n(\tau,\gamma,\varphi)$ if the following conditions are satisfied:*

$$1 + \frac{1}{\tau}\left[(D^nf(z))' + \gamma z(D^nf(z))'' - 1\right] \prec \varphi(z)$$

$(0 \leq \gamma \leq 1,\ \tau \in \mathbb{C}/\{0\},\ n \in \mathbb{N},\ z \in \mathcal{U})$ *and*

$$1 + \frac{1}{\tau}\left[(D^ng(w))' + \gamma w(D^ng(w))'' - 1\right] \prec \varphi(w)$$

$(0 \leq \gamma \leq 1,\ \tau \in \mathbb{C}/\{0\},\ n \in \mathbb{N},\ w \in \mathcal{U})$, *where the function g is given by $g(w) = f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + ...$ and D^n is the Salagean differential operator.*

In this paper, we obtain the estimates on the coefficients $|a_2|$ and $|a_3|$ for $\Sigma^n(\tau,\gamma,\varphi)$ as well as its special classes.

Firstly, in order to derive our main results, we need the following lemma.

Lemma 1.1. [8] *Let $p(z) = 1 + c_1 z + c_2 z^2 + ... \in P$, where P is the family of all functions p, analytic in $\mathcal{U}$, for which $\operatorname{Re} p(z) > 0$ $(z \in \mathcal{U})$. Then*

$$|c_n| \leq 2; \ n = 1, 2, 3, ...$$

2. INITIAL COEFFICIENTS FOR THE CLASS $\Sigma^n(\tau, \gamma, \varphi)$

Theorem 2.1. *Let $f(z) \in \Sigma^n(\tau, \gamma, \varphi)$ be of the form (1.1). Then*

$$|a_2| \leq \frac{|\tau| \sqrt{B_1^3}}{\sqrt{\left|3^{n+1}\tau B_1^2 (1+2\gamma) + 4^{n+1}(1+\gamma)^2 (B_1 - B_2)\right|}} \tag{2.1}$$

and

$$|a_3| \leq B_1 |\tau| \left(\frac{B_1 |\tau|}{4^{n+1}(1+\gamma)^2} + \frac{1}{3^{n+1}(1+2\gamma)} \right). \tag{2.2}$$

Proof. Since $f \in \Sigma^n(\tau, \gamma, \varphi)$, there exist two analytic functions $u, v : \mathcal{U} \to \mathcal{U}$, with $u(0) = v(0) = 0$, such that

$$1 + \frac{1}{\tau}\left[(D^n f(z))' + \gamma z (D^n f(z))'' - 1\right] = \varphi(u(z)) \quad (z \in \mathcal{U}) \tag{2.3}$$

and

$$1 + \frac{1}{\tau}\left[(D^n g(w))' + \gamma w (D^n g(w))'' - 1\right] = \varphi(v(w)) \quad (w \in \mathcal{U}). \tag{2.4}$$

Define the function p and q as following:

$$p(z) = \frac{1+u(z)}{1-u(z)} = 1 + c_1 z + c_2 z^2 + c_3 z^3 + ...$$

and

$$q(w) = \frac{1+v(w)}{1-v(w)} = 1 + b_1 w + b_2 w^2 + b_3 w^3 + ...$$

or equivalently,

$$u(z) = \frac{p(z)-1}{p(z)+1} = \frac{c_1}{2} z + \frac{1}{2}\left(c_2 - \frac{c_1^2}{2}\right) z^2 + \frac{1}{2}\left(c_3 + \frac{c_1}{2}\left(\frac{c_1^2}{2} - c_2\right) - \frac{c_1 c_2}{2}\right) z^3 ... \tag{2.5}$$

and

$$v(w) = \frac{q(w)-1}{q(w)+1} = \frac{b_1}{2} w + \frac{1}{2}\left(b_2 - \frac{b_1^2}{2}\right) w^2 + \frac{1}{2}\left(b_3 + \frac{b_1}{2}\left(\frac{b_1^2}{2} - b_2\right) - \frac{b_1 b_2}{2}\right) w^3 .... \tag{2.6}$$

If we use (2.5) and (2.6) in (2.3) and (2.4) along with (1.3), we have

$$1 + \frac{1}{\tau}\left[(D^n f(z))' + \gamma z (D^n f(z))'' - 1\right]$$
$$= 1 + \frac{1}{2} B_1 c_1 z + \left[\frac{1}{2} B_1 \left(c_2 - \frac{c_1^2}{2}\right) + \frac{1}{4} B_2 c_1^2\right] z^2 + ... \tag{2.7}$$

and

$$1+\frac{1}{\tau}\left[\left(D^{n}g\left(w\right)\right)'+\gamma w\left(D^{n}g\left(w\right)\right)''-1\right]$$
$$=1+\frac{1}{2}B_{1}b_{1}w+\left[\frac{1}{2}B_{1}\left(b_{2}-\frac{b_{1}^{2}}{2}\right)+\frac{1}{4}B_{2}b_{1}^{2}\right]w^{2}+.... \tag{2.8}$$

It follows from (2.7) and (2.8) that

$$\frac{\left(1+\gamma\right)2^{n+1}a_{2}}{\tau}=\frac{1}{2}B_{1}c_{1} \tag{2.9}$$

$$\frac{3^{n+1}\left(1+2\gamma\right)a_{3}}{\tau}=\frac{1}{2}B_{1}\left(c_{2}-\frac{c_{1}^{2}}{2}\right)+\frac{1}{4}B_{2}c_{1}^{2} \tag{2.10}$$

and

$$-\frac{\left(1+\gamma\right)2^{n+1}a_{2}}{\tau}=\frac{1}{2}B_{1}b_{1} \tag{2.11}$$

$$\frac{3^{n+1}\left(1+2\gamma\right)\left(2a_{2}^{2}-a_{3}\right)}{\tau}=\frac{1}{2}B_{1}\left(b_{2}-\frac{b_{1}^{2}}{2}\right)+\frac{1}{4}B_{2}b_{1}^{2}. \tag{2.12}$$

From (2.9) and (2.11) we obtain

$$c_{1}=-b_{1} \tag{2.13}$$

By adding (2.10) to (2.12) and combining this with (2.9) and (2.11), we get

$$a_{2}^{2}=\frac{\tau^{2}B_{1}^{3}\left(b_{2}+c_{2}\right)}{4\left[3^{n+1}\tau B_{1}^{2}\left(1+2\gamma\right)+4^{n+1}\left(1+\gamma\right)^{2}\left(B_{1}-B_{2}\right)\right]}. \tag{2.14}$$

Subtracting (2.10) from (2.12), if we use (2.9) and applying (2.13), we have

$$a_{3}=\frac{\tau^{2}B_{1}^{2}b_{1}^{2}}{2^{2n+4}\left(1+\gamma\right)^{2}}+\frac{\tau B_{1}\left(c_{2}-b_{2}\right)}{4.3^{n+1}\left(1+2\gamma\right)}. \tag{2.15}$$

Finally, in view of Lemma 1.1, we get results (2.1) to (2.2) asserted by the Theorem 2.1. □

3. COROLLARIES AND CONSEQUENCES

i) If we set

$$\tau=e^{i\beta}\cos\beta\quad\left(-\frac{\pi}{2}<\beta<\frac{\pi}{2}\right)$$

and

$$\varphi(z)=\frac{1+(1-2\kappa)z}{1-z}=1+2(1-\kappa)z+2(1-\kappa)z^{2}+...\quad(0\leq\kappa<1)$$

which gives $B_1=B_2=2(1-\kappa)$, in Theorem 2.1, we can have the following corollary.

Corollary 3.1. *Let* $f(z) \in \Sigma^n \left(e^{i\beta}\cos\beta\ , \gamma, \frac{1+(1-2\kappa)z}{1-z}\right)$ *be of the form (1.1). Then*

$$|a_2| \leq \sqrt{\frac{2\,(1-\kappa)}{3^{n+1}(1+2\gamma)}\cos\beta} \tag{3.1}$$

and

$$|a_3| \leq 2\,(1-\kappa)\left(\frac{(1-\kappa)\cos\beta}{2^{2n+1}\,(1+\gamma)^2} + \frac{1}{3^{n+1}\,(1+2\gamma)}\right)\cos\beta. \tag{3.2}$$

Remark 3.2. *For* $\gamma = 0$, *Corollary 3.1 simplifies to the following form.*

Corollary 3.3. *Let* $f(z) \in \Sigma^n \left(e^{i\beta}\cos\beta\ , 0, \frac{1+(1-2\kappa)z}{1-z}\right)$ *be of the form (1.1). Then*

$$|a_2| \leq \sqrt{\frac{2\,(1-\kappa)}{3^{n+1}}\cos\beta} \tag{3.3}$$

and

$$|a_3| \leq 2\,(1-\kappa)\left(\frac{(1-\kappa)\cos\beta}{2^{2n+1}} + \frac{1}{3^{n+1}}\right)\cos\beta. \tag{3.4}$$

ii) If we set $\tau = 1$ and

$$\varphi(z) = \left(\frac{1+z}{1-z}\right)^{\alpha} = 1 + 2\alpha z + 2\alpha^2 z^2 + \ldots \quad (0 < \alpha \leq 1)$$

which gives $B_1 = 2\alpha$, $B_2 = 2\alpha^2$, in Theorem 2.1, we can obtain the following corollary.

Corollary 3.4. *Let* $f(z) \in \Sigma^n \left(1, \gamma, \left(\frac{1+z}{1-z}\right)^{\alpha}\right)$ *be of the form (1.1). Then*

$$|a_2| \leq \alpha\sqrt{\frac{2}{3^{n+1}(1+2\gamma)\,\alpha + 2^{2n+1}\,(1+\gamma)^2\,(1-\alpha)}} \tag{3.5}$$

and

$$|a_3| \leq \left(\frac{\alpha^2}{4^n\,(1+\gamma)^2} + \frac{2\alpha}{3^{n+1}\,(1+2\gamma)}\right). \tag{3.6}$$

Remark 3.5. *In its special case when* $\gamma = 0$ *in Corollary 3.4, we can get the following corollary.*

Corollary 3.6. *Let* $f(z) \in \Sigma^n \left(1, 0, \left(\frac{1+z}{1-z}\right)^{\alpha}\right)$ *be of the form (1.1). Then*

$$|a_2| \leq \alpha\sqrt{\frac{2}{\alpha 3^{n+1} + 2^{2n+1}\,(1-\alpha)}} \tag{3.7}$$

and

$$|a_3| \leq \left(\frac{\alpha^2}{4^n} + \frac{2\alpha}{3^{n+1}}\right). \tag{3.8}$$

Remark 3.7. **i:** *Putting $n = 0$ in Theorem 2.1, we obtain the corresponding result given earlier by Deniz* [7] *(also Srivastava and Bansal* [21]*).*

ii: *Putting $\tau = 1$, $\gamma = 0$, $n = 0$ in Theorem 2.1, we obtain the corresponding result given earlier by Ali et al* [1].

iii: *Putting $\beta = 0$, $n = 0$ in Corollary 3.3 and $\gamma = 0$, $n = 0$ in Corollary 3.4, we obtain the corresponding result given earlier by Srivastava et al* [19].

References

[1] R. M. Ali, S. K. Lee, V. Ravichandran and S. Supramaniam, Coefficient estimates for bi-univalent Ma-Minda starlike and convex functions, Appl. Math. Lett. 25 (3) (2012), 344–351.

[2] D. A. Brannan, J.G. Clunie (Eds.), Aspects of Contemporary Complex Analysis (Proceedings of the NATO Advanced Study Institute held at the University of Durham, Durham; July 1 20, 1979), Academic Press, New York and London, 1980.

[3] D. A. Brannan and T.S. Taha, On some classes of bi-univalent functions, in: S.M. Mazhar, A. Hamoui, N.S. Faour (Eds.), Math. Anal. and Appl., Kuwait; February 18–21, 1985, in: KFAS Proceedings Series, vol. 3, Pergamon Press, Elsevier Science Limited, Oxford, 1988, pp. 53–60. see also Studia Univ. Babeş-Bolyai Math. 31 (2) (1986), 70–77.

[4] S. Bulut, Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions, C. R. Acad. Sci. Paris, Ser. I, 352 (6) (2014), pp. 479–484.

[5] S. Bulut, N. Magesh and V. K. Balaji, Faber polynomial coefficient estimates for certain subclasses of meromorphic bi-univalent functions, Comptes Rendus Mathematique 353(2) (2015), 113-116.

[6] M. Çağlar, H. Orhan and N. Yağmur, Coefficient bounds for new subclasses of bi-univalent functions, Filomat 27(7) (2013), 1165-1171.

[7] E. Deniz, Certain subclasses of bi-univalent functions satisfying subordinate conditions, J. Class. Anal. 2(1) (2013), 49–60.

[8] P. L. Duren, Univalent functions,Grundlehren der Mathematischen Wissenschaften, 259, Springer, New York, 1983.

[9] B. A. Frasin, M.K. Aouf, New subclasses of bi-univalent functions, Appl. Math. Lett. 24 (2011), 1569-1573.

[10] S. G. Hamidi, J. M. Jahangiri, Faber polynomial coefficient estimates for analytic bi-close-to-convex functions, C. R. Acad. Sci. Paris, Ser. I 352 (2014), 17–20.

[11] A. W. Kedzierawski, Some remarks on bi-univalent functions, Ann. Univ.Mariae Curie-Skłodowska Sect. A 39 (1985), 77–81 (1988).

[12] S. Sivaprasad Kumar, V. Kumar and V. Ravichandran, Estimates for the initial coefficients of bi-univalent functions, Tamsui Oxford J. Inform. Math. Sci. 29(4) (2013), 487-504.

[13] M. Lewin, On a coefficient problem for bi-univalent functions, Proc. Amer. Math. Soc. 18 (1967), 63–68.

[14] A. K. Mishra and S. Barık, Estimates for the initial coefficients of bi-univalent $\lambda-$convex analytic functions in the unit disc, Journal of Classical Analysis, 7(1) (2015), 73-81.

[15] E. Netanyahu, The minimal distance of the image boundary from the origin and the second coefficient of a univalent function in $|z| < 1$, Arch. Rational Mech. Anal. 32 (1969), 100-112.

[16] H. Orhan, N. Magesh and V. K. Balaji, Initial coefficient bounds for a general class of bi-univalent functions, Filomat 29(6) (2015), 1259–1267.

[17] C. Ramachandran, R. Ambrose Prabhu and N. Magesh, Initial coefficient estimates for certain subclasses of bi-univalent functions of Ma-Minda type, Applied Mathematical Sciences, 9(47) (2015), 2299-2308.

[18] G. S. Salagean, Subclasses of univalent functions, Complex analysis - Proc. 5th Rom.-Finn. Semin., Bucharest 1981, Part 1, Lect. Notes Math., 1013 (1983), 362–372.

[19] H. M. Srivastava, A. K. Mishra, P. Gochhayat, Certain subclasses of analytic and bi-univalent functions, Appl. Math. Lett. 23 (2010), 1188–1192.
[20] H. M. Srivastava, S. Bulut, M. Çağlar, N. Yağmur, Coefficient estimates for a general subclass of analytic and bi-univalent functions, Filomat 27(5) (2013), 831-842.
[21] H. M. Srivastava and D. Bansal, Coefficient estimates for a subclass of analytic and bi-univalent functions, Journal of the Egyptian Mathematical Society 23(2) (2015), 242-246.
[22] Y. Sun, Y. P. Jiang and A. Rasila, Coefficient estimates for certain subclasses of analytic and bi-univalent functions, Filomat 29(2) (2015), 351-360.
[23] D. L. Tan, Coefficient estimates for bi-univalent functions, Chinese Ann. Math. Ser. A 5 (5) (1984), 559–568.
[24] P. Zaprawa, Estimates of initial coefficients for bi-univalent functions, Abstr. Appl. Anal., 2014, Article ID 357480, 6 pages.

Current address: Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars, Turkey.
E-mail address, Murat Çağlar: mcaglar25@gmail.com (corresponding author)
Current address: Department of Mathematics, Faculty of Science and Letters, Kafkas University, Kars, Turkey.
E-mail address, Erhan Deniz: edeniz36@gmail.com